

97/2002

THIRAM 320

Flowable Fungicide

A seed treatment for control of
verticillium wilt of alfalfa.

NET CONTENTS: 20 litres

AGRICULTURAL

GUARANTEE: THIRAM.....32.4%

REGISTRATION NO. 16189
PEST CONTROL PRODUCTS ACT

READ THE LABEL BEFORE USING

DANGER POISON

UNIROYAL CHEMICAL LTD.
25 Erb Street
Elmira, Ontario
N3B 3A3

24 Hour Emergency Phone: (519) 744-3060

DIRECTIONS FOR USE:

SEED TREATMENT: For control of verticillium wilt of alfalfa, apply 720 mL of THIRAM 320 per 100 kilograms of seed. Prior to application, THIRAM 320 flowable fungicide should be thoroughly agitated to insure uniform mixing of the product.

DO NOT DILUTE THIRAM 320 with additional water.

Add THIRAM 320 to the seed and mix well for three minutes in treating equipment. Seed may be treated in cement mixer or commercially available seed treating equipment.

Thorough coverage of the seed is required.

Do not store treated seed for more than 5 months.

USE RESTRICTIONS:

1. Label treated seed "DO NOT USE FOR FOOD OR FEED. THIS SEED HAS BEEN TREATED WITH THIRAM".
2. Store in a cool, dry, ventilated place away from feeds and foods.
3. Do not contaminate food or feed.
4. Keep away from heat, fire and sparks.
5. Treated seed may be harmful to birds if ingested.
6. Do not contaminate any body of water.

SAFETY PRECAUTIONS:

1. KEEP OUT OF REACH OF CHILDREN.
2. May irritate eyes, nose, throat and skin.
3. May cause allergenic eczema in sensitive individuals.
4. Avoid contact with skin, eyes and clothing.
5. Wash thoroughly after handling and before eating, drinking or smoking.
6. Wash contaminated clothing with soap and hot water before reuse.
7. Do not consume alcoholic beverages for 24 hours before or after working with THIRAM 320.
8. Avoid breathing vapours.
9. When treating, augering or handling treated seed, work in a well ventilated area and wear a suitable mask, goggles and gloves.

SYMPTOMS OF POISONING: Skin contact may cause irritation and dermatitis. Alcohol intake within 24 hours of contact with THIRAM may cause flushing, sweating, headache and nausea.

FIRST AID:

IF SWALLOWED: Induce vomiting by giving the patient syrup of Ipecac (adult or child 12 years or older 30 mL; child under 12, 15 mL) followed by water to enhance vomiting. Keep patient quiet. Seek medical attention IMMEDIATELY and bring this labelled container with you.

IF IN EYES: Wash immediately with running water for 15 minutes. Obtain medical attention.

IF ON THE SKIN: Wash with warm water and soap for 15 minutes.

TOXICOLOGICAL INFORMATION: There is no specific antidote. If swallowed, induce vomiting with SYRUP OF IPECAC or INTUBATE the stomach. If reaction with alcohol occurs treat as disulfiram-alcohol reaction.

DISPOSAL:

1. Follow provincial instructions for any required additional cleaning of the container prior to its disposal.
2. Make the empty container unsuitable for further use.
3. Dispose of the container in accordance with provincial requirements.
4. For information on the disposal of unused, unwanted product and the cleanup of spills, contact the Provincial Regulatory Agency or the Manufacturer.

NOTICE TO USER: This control product is to be used only in accordance with the directions on this label. It is an offence under the Pest Control Products Act to use a control product under unsafe conditions.

NOTICE TO BUYER: Seller's guarantee shall be limited to the terms set out on the label and subject thereto, the buyer assumes the risk to persons or property arising from the use or handling of this product and accepts the product on that condition.

This label transcript service is offered by the Pest

Management Regulatory Agency to provide efficient searching for label information. This service and this information do not replace the official hard-copy label. The PMRA does not provide any guarantee or assurance that the information obtained through this service is accurate, current or correct, and is therefore not liable for any loss resulting, directly or indirectly, from reliance upon this service.

+))