

(Booklet)

Group	4	Herbicide
-------	---	-----------

Tracker®
LIQUID HERBICIDE FOR CONTROL OF BROADLEAF WEEDS

AGRICULTURAL

GUARANTEE: MCPA (present as amine salts)..... 275 g/L
Mecoprop (present as amine salts)..... 62.5 g/L
Dicamba (present as amine salts)..... 62.5 g/L

REGISTRATION NO.: 25605

PEST CONTROL PRODUCTS ACT



CAUTION

POISON

NET CONTENTS: 10 L

**READ THE LABEL AND ATTACHED BROCHURE BEFORE USING.
PROTECT FROM FREEZING.**

INTERPROVINCIAL COOPERATIVE LIMITED
P.O. BOX 1050
SASKATOON, SASKATCHEWAN S7K 3M9

RC000-0904

GENERAL INFORMATION:

For systemic control of many broadleaf weeds in: Wheat (spring, durum and winter), Barley, Oats, stubble fields, Summerfallow, and Annual Canary Grass.

Note: Do not underseed to legumes.

WEEDS CONTROLLED:

Best results are obtained when weeds are in the 2-3 leaf stage and actively growing. Less than satisfactory control may result if weeds are not actively growing. Buckwheat (wild, tartary, volunteer cultivated), Cleavers¹, Common Ragweed, Corn Spurry, Cow Cockle, Flixweed, Green Smartweed, Hempnettle², Knotweed, Kochia, Lady's-thumb, Lamb's-quarters, Mustards (wild, ball, tall, wormseed, yellow), Night Flowering Catchfly, Pigweeds (prostrate, redroot), Russian Thistle³, Shepherd's Purse, Annual Sow Thistle, Stinkweed, Volunteer Rapeseed, Volunteer Sunflower.

Top Growth Control Only:

Canada Thistle, Field Bindweed, Hedge Bindweed, Perennial Sow Thistle

Note:

¹Cleavers: Spray in 1-2 whorl stage

²Hempnettle: Spray before second pair of true leaves

³Russian Thistle: Spray when less than 5 cm in height

TIMING OF APPLICATION:

Crop	Crop State
Wheat (spring, durum) Oats Annual Canary Grass	2 to 5-leaf
Winter Wheat	For winter wheat, apply in the Spring before the crop is more than 30 cm high (top leaf extended). Winter annual weeds should be sprayed as soon as new growth starts.
Barley	2 to 4-leaf

Note: Application made later than the recommended timing may result in flattening of the crop and shortening of the straw (particularly under stress conditions).

Application information:

Rate 1.0 to 1.5 litres per hectare

Use the 1.5 L/ha rate:

- a) when weeds are not actively growing due to extended periods of hot and dry or cold and wet weather prior to or following application.
- b) when weeds are beyond the 3-leaf stage
- c) for heavy weed infestations.
- d) for top growth control of Canada Thistle, Perennial Sow Thistle, Field Bindweed, Hedge Bindweed and for control of Cleavers.
- e) For control of overwintering Flixweed, Shepherd's-Purse and stinkweed.

Avoid overlapping. Shut off spray boom while starting, turning, slowing and stopping, otherwise crop injury may result.

BROADLEAF WEEDS, WILD OATS, GREEN AND YELLOW FOXTAIL CONTROL IN SPRING WHEAT ONLY IN THE PRAIRIE PROVINCES AND THE PEACE RIVER REGION OF BRITISH COLUMBIA.

To control broadleaf weeds, wild oats and yellow foxtail in spring wheat only in the following tank mix:

	Broadleaf Weeds and Wild Oats	Green Foxtail Wild Oats Yellow Foxtail and Broadleaf Weeds
TRACKER 1.0 - 1.5 L/ha+	230 mL/ha of HORIZON 240 EC + 0.8% volume/volume of SCORE (800 mL of SCORE in 100 L of water per Hectare)	290 mL/ha of HORIZON 240 EC + 1.0% volume/volume of SCORE (1 L of SCORE with 100 L of water per hectare)

Note:

1. Do not use **TRACKER/HORIZON** tank mix on barley or on oats.
2. Consult tank mix partner label for detailed instruction and precautions.
3. Apply when spring wheat is in the 2-5 leaf stage.
4. DO NOT APPLY THIS TANK MIX BY AIR.
5. Use a minimum of 100 L of water per hectare.
6. Observe a minimum interval to harvest of 60 days after treatment.
7. Observe a minimum of 7 days before grazing livestock on crops treated with the tank mix.
8. Apply when broadleaf weeds are in the 2-3 leaf stage, when wild oats are in the 1-6 leaf stage and when foxtails are in the 1-5 leaf stage.
9. Do not contaminate streams, rivers, lakes, ponds, wetlands, sloughs, irrigation waters, vegetated ditch banks and other water used by livestock or for domestic purposes. Do not apply within 15 metres of a wetland area or any of the aforementioned bodies of water.

CHICKWEED CONTROL:

To control Chickweed and other hard-to-kill broadleaf weeds in wheat and barley, use one of the following tank mixes:

		TANK MIX PARTNER	RATE
		SENCOR® 500 F**	275-425 mL/ha
TRACKER®			
1.0-1.5 L/ha	+	LOROX® L	425 mL/ha

Consult the tank mix partner label for detailed instructions and precautions.

** For Western Canada only: Application of this tank mix at later crop stages may result in injury if crop is under stress.

CANADA THISTLE:

Program for gradual eradication of Canada Thistle:

Year 1: Summerfallow	Apply 2 L/ha when Canada Thistle is in early bud stage. Do not cultivate for a minimum of 4 weeks after application.
Or: Post Harvest (Stubble)	Apply 2 L/ha to Canada Thistle re-growth. Treatment at the 15-20 cm stage of actively growing Canada Thistle will provide the best results. Do not apply within 2 weeks of killing frost.

Year 2: In Crop	Apply 1.5 litres per hectare when Canada Thistle is young and actively growing.
-----------------	---

Note: Repeating this program will gradually eliminate Canada Thistle.

MIXING AND SPRAYING INSTRUCTIONS:

GROUND APPLICATION

Spray pressure: 200-300 kPa

Spray volume: minimum of 100 litres of water per hectare

Use 50 mesh screen and filters

Note: DO NOT add an acidifier to the spray mixture.

1. Fill the sprayer $\frac{1}{2}$ full with water. Start gentle agitation.
2. Add tank mix partner, if applicable, and agitate thoroughly.
3. Fill tank $\frac{3}{4}$ full with water and add **TRACKER**.
4. Complete filling of sprayer and maintain gentle agitation while mixing and spraying.
5. Do not allow contents to stand without agitation.

AERIAL APPLICATION

· Boom pressure: 235 kPa or less

· Spray volume: minimum of 30 litres of water per hectare

· Use 50 mesh screens and filters

· Avoid placing nozzles where spray will enter wing tip vortices.

Note: DO NOT add an acidifier to the spray mixture.

1. Fill the mixing tank $\frac{1}{2}$ full with water. Start gentle agitation.
2. Add tank mix partner, if applicable, and agitate thoroughly.
3. Fill mixing tank $\frac{3}{4}$ full with water and add **TRACKER**.
4. Complete filling of mixing tank and maintain gentle agitation.
5. Fill aircraft spray tank and maintain gentle agitation while

- spraying.
6. Do not let contents stand without agitation.

AERIAL APPLICATION USE PRECAUTIONS

- Coarse sprays are less likely to drift. Use only nozzles or nozzle configurations which minimize the production of fine spray drops. Do not angle nozzles forward into the airstream and do not increase spray volume by increasing nozzle pressure. When spraying avoid combinations of pressure and nozzle type that will result in fine particles (mist) which are more likely to drift. A spray thickening agent or drift retardant may be used with this product to aid in reducing spray drift.
- Avoid direct applications to any body of water. Do not contaminate water through spray drift or by cleaning of equipment or disposal of wastes.
- Do not use human flaggers. Avoid spray drift. Apply only when there is little or no hazard from spray drift. Small quantities of the spray, which may not be visible, may seriously injure susceptible crops and damage sensitive non-target habitat. A method, such as balloons or a continuous smoke column at or near the spray site or a smoke generator on the spray equipment, must be used to detect air movement, lapse conditions or temperature inversions (stable air). If the smoke develops into layers or indicates a potential for hazardous spray drift, do not spray.
- Do not apply TRACKER directly to, or otherwise permit it to come into direct contact with desirable crops or other desirable broadleaf plants or non-target species and do not permit spray mists to drift onto them.
- **BUFFER ZONE:** Observe the provincial buffer zones between treatment areas and any aquatic systems or significant wildlife habitat.

PRECAUTIONS:

KEEP OUT OF REACH OF CHILDREN. Do not reuse containers. Do not allow spray drift to contact plants other than those being sprayed. Do not contaminate streams, lakes, ponds, and irrigation water, water used for livestock, or domestic purpose. Sprayers which have contained **TRACKER** should not be used to apply other pesticides to MCPA or 2,4-D susceptible crops.

If MCPA or 2,4-D susceptible plants are growing within a distance of 400 meters from the area being sprayed, spray only when there is a light breeze away from the susceptible plants. Spray drifting can be reduced by using high volume sprays under low pressure, coarse sprays and drop nozzles. Susceptible crops include grape, tomato, tobacco, sugar beet, sunflower, beans, soybean, turnip, cauliflower and cabbage as well as many other field, vegetable, ornamental and fruit crops. If **TRACKER** is applied when crop is under stress, shortening of straw may occur, but yield will not be affected. Protect from freezing.

If this pest control product is to be used on a commodity that may be exported to the U.S. and you require information on acceptable residue levels in the U.S., contact 1-866-375-4648 or www.cropro.org.

RESISTANCE MANAGEMENT RECOMMENDATIONS: For resistance management **TRACKER** is a group 4 herbicide. Any weed population may contain or develop plants naturally resistant to **TRACKER** and other Group 4 herbicides. The resistant biotypes may dominate the weed population if these herbicides are used repeatedly in the same field. Other resistance mechanisms that are not linked to site of action, but specific for individual chemicals, such as enhanced metabolism, may also exist. Appropriate resistance-management strategies should be followed.

To delay herbicide resistance: Where possible, rotate the use of **TRACKER** or other Group 4 herbicides with different herbicide groups that control the same weeds in a field. Use tank mixtures with herbicides from a different group when such use is permitted. Herbicide use should be based on an IPM program that includes scouting, historical information related to herbicide use and crop rotation and considers tillage (or other mechanical), cultural, biological and other chemical control practices. Monitor treated weed populations for resistance development. Prevent movement or resistant weed seeds to other fields by cleaning harvesting and tillage equipment and planting clean seed. Contact your local extension specialist or certified crop advisors for any additional pesticide resistance-management and/or integrated weed-management recommendations for specific crops and weed biotypes. For further information or to report suspected resistance, contact Interprovincial Cooperative Ltd. at 1-204-233-3461.

CAUTION:

Do not graze or harvest for livestock feed within seven (7) days of application.

TOXICOLOGICAL INFORMATION: No specific antidote. Treatment of any systemic intoxication should be primarily symptomatic and supportive.

FIRST AID:

If on skin, wash with soap and water. **If in eyes**, flush with clean water for 15 minutes. If irritation persists, get medical attention. **If swallowed**, induce vomiting. Obtain prompt medical attention or contact a poison control centre IMMEDIATELY.

Take container, label or product name and Pest Control Product Registration Number with you when seeking medical attention.

DISPOSAL:

Do not reuse this container for any purpose. This is a recyclable container, and is to be disposed of at a container collection site. Contact your local distributor/dealer or municipality for the location of the nearest collection site. Before taking the container to the collection site:

1. Triple or pressure rinse the empty container. Add the rinsings to the spray mixture in the tank.
2. Make the empty, rinsed container unsuitable for further use.

If there is no container collection site in your area, dispose of the container in accordance with provincial requirements.

For information on disposal of unused, unwanted product, contact the manufacturer or the provincial regulatory agency. Contact the manufacturer and the provincial regulatory agency in case of a spill, and for clean-up of spills.

NOTICE TO USER:

This control product is to be used only in accordance with the directions on this label. It is an offense under the *Pest Control Products Act* to use a control product under unsafe conditions.

NOTICE TO BUYER:

Seller's guarantee shall be limited to the terms set out on the label and, subject thereto, the buyer assumes the risk to persons or property arising from the use or handling of this product and accepts the product on that condition.

All product names are registered trademarks of their respective companies.

(Base Label)

Group	4	Herbicide
-------	---	-----------

Tracker®

LIQUID HERBICIDE FOR CONTROL OF BROADLEAF WEEDS

AGRICULTURAL

GUARANTEE: MCPA (present as amine salts)..... 275 g/L
Mecoprop (present as amine salts)..... 62.5 g/L
Dicamba (present as amine salts)..... 62.5 g/L

REGISTRATION NO.: 25605

PEST CONTROL PRODUCTS ACT



CAUTION

POISON

NET CONTENTS: 10 L

**READ THE LABEL AND ATTACHED BROCHURE BEFORE USING.
PROTECT FROM FREEZING.**

INTERPROVINCIAL COOPERATIVE LIMITED
P.O. BOX 1050
SASKATOON, SASKATCHEWAN S7K 3M9

RC000-0904

(Back Panel)

TRACKER®

PRECAUTIONS:

KEEP OUT OF REACH OF CHILDREN. Do not reuse containers. Do not allow spray drift to contact plants other than those being sprayed. Do not contaminate streams, lakes, ponds, and irrigation water, water used for livestock, or domestic purpose. Sprayers which have contained **TRACKER** should not be used to apply other pesticides to MCPA or 2,4-D susceptible crops.

If 2,4-D or MCPA susceptible plants are growing within a distance of 400 metres from the area being sprayed, spray only when there is a light breeze away from the susceptible plants. Spray drifting can be reduced by using high volume sprays under low pressure, coarse sprays and drop nozzles. Susceptible crops include grape, tomato, tobacco, sugar beet, sunflower, beans, soybean, turnip, cauliflower and cabbage as well as many other field, vegetable, ornamental and fruit crops. If **TRACKER** is applied when crop is under stress, shortening of straw may occur, but yield will not be affected. Protect from freezing.

If this pest control product is to be used on a commodity that may be exported to the U.S. and you require information on acceptable residue levels in U.S., contact 1-866-375-4648 or www.croplife.ca.

CAUTION:

Do not graze or harvest for livestock feed within seven (7) days of application.

FIRST AID:

If on skin, wash with soap and water. **If in eyes**, flush with clean water for 15 minutes. If irritation persists, get medical attention. **If swallowed**, induce vomiting. Obtain prompt medical attention or contact a poison control centre IMMEDIATELY.

Take container, label or product name and Pest Control Product Registration Number with you when seeking medical attention.

DISPOSAL:

Do not reuse this container for any purpose. This is a recyclable container, and is to be disposed of at a container collection site. Contact your local distributor/dealer or municipality for the location of the nearest collection site. Before taking the container to the collection site:

1. Triple or pressure rinse the empty container. Add the rinsings to the spray mixture in the tank.
2. Make the empty, rinsed container unsuitable for further use.

If there is no container collection site in your area, dispose of the container in accordance with provincial requirements.

For information on disposal of unused, unwanted product, contact the manufacturer or the provincial regulatory agency. Contact the manufacturer and the provincial regulatory agency in case of a spill, and for clean-up of spills.

NOTICE TO USER:

This control product is to be used only in accordance with the directions on this label. It is an offense under the *Pest Control Products Act* to use a control product under unsafe conditions.

NOTICE TO BUYER:

Seller's guarantee shall be limited to the terms set out on the label and, subject thereto, the buyer assumes the risk to persons or property arising from the use or handling of this product and accepts the product on that condition.

®T.M. Reg'd/Marque deposee

RC000-0904

This label transcript service is offered by the Pest Management Regulatory Agency to provide efficient searching for label information. This service and this information do not replace the official hard-copy label. The PMRA does not provide any guarantee or assurance that the information obtained through this service is accurate, current or correct, and is therefore not liable for any loss resulting, directly or indirectly, from reliance upon this service.