((LABEL))

RAMPEX

OUTDOOR CRAWLING INSECT KILLER

REG. NO. 25180 P.C.P. ACT

GUARANTEE: Chlorpyrifos (Dursban (R)) 50 g/L

DOMESTIC

KEEP OUT OF REACH OF CHILDREN

DANGER * POISON

READ THE LABEL AND CARTON BEFORE USING

500 mT₁

REFER TO AND RETAIN CARTON FOR "DIRECTIONS FOR USE".

PRECAUTIONS: KEEP OUT OF REACH OF CHILDREN. May be fatal if swallowed. May be absorbed through the skin. May be injurious to eyes and skin. Wash hands and exposed skin after spraying, also any contaminated clothing. Avoid breathing spray mist. Do not permit pets to contact sprayed areas until dry. Do not contaminate food, feed or water; product is poisonous to fish. Re-close container tightly after use and store away from heat.

FIRST AID: Product contains petroleum distillates. If swallowed DO NOT induce vomiting but rush patient to hospital, taking this labelled container with you. If in eyes, flush with plenty of water for 15 minutes, and get medical aid. If on skin wash thoroughly with soap and water.

TOXICOLOGICAL INFORMATION: Chlorpyrifos is a cholinesterase inhibitor; atropine by injection only is antidotal.

DISPOSAL: Dispose of empty container with household garbage. Do not re-use container.

(R) Dursban is a registered trademark of DowElanco

((CARTON))

RAMPEX

OUTDOOR CRAWLING INSECT KILLER

REG. NO. 25180 P.C.P. ACT

GUARANTEE: Chlorpyrifos 50 g/L (Dursban (R))

DOMESTIC

KEEP OUT OF REACH OF CHILDREN

DANGER * POISON

READ THE LABEL AND CARTON BEFORE USING

500 mL

Maheu & Maheu Inc.
710, rue Bouvier, #195
 Québec (Québec)
 G2J 1C2
Tél.: (418) 623-8000
1-800-463-2186

5 mL= 1 Teaspoonful

15 mL= 1 Tablespoonful

DIRECTIONS FOR USE:

TURF AND LAWNS: Make no more than 2 applications per season. To control WHITE GRUBS and the grubs of the EUROPEAN CHAFER and the JAPANESE BEETLE mix 200~mL in 5~L of water and apply evenly over 50~mL

m2. Spray in late July or August when grubs are small and are feeding on the grass roots just below soil surface. IMMEDIATELY AFTER SPRAYING, WATER THE LAWNS THOROUGHLY. To control CHINCH BUGS, ANTS and MOSQUITOES, SOD WEBWORMS, larvae of (lawn moths), mix 200 mL in 10 L of water and apply to 100 m2 of lawn area (a watering can is suitable for this treatment). Apply when these insects first appear and repeat when needed. The lawn surface should be moist before making the treatment. For Sod Webworms delay watering or mowing for 12 to 24 hours after treatment. To control CRANE FLY LARVAE (leatherjackets) use 200 mL in 10 L of water for each 100 m2. Apply as a drenching spray in water in the late fall after the flight of adult crane flies has ceased for the year.

ORNAMENTAL GARDENS: Make no more than 2 treatments per season. To control ANTS, CRICKETS, EARWIGS, SOWBUGS, SPIDERS, CUTWORMS, WHITE GRUBS, EUROPEAN CHAFER LARVAE, and JAPANESE BEETLE LARVAE. Spray soil around plants to give good coverage. Use 200 mL in 10 L of water and apply to 100 m2 of soil.

For the following foliar insect pests; aphids, ash and lilac borers, mealybugs, mites, scales, spittlebugs, thrips and whiteflies mix 20 mL per L of water and spray to give uniform coverage. Do not use on roses, camelias, poinsettias, geraniums, oxalis, coleus or variegated ivy.

NATIVE ELM BARK BEETLE: Make only one application per season. To prevent the adult beetle from overwintering in uninfected trees and to reduce beetle populations in disease free areas, mix 250 mL in 2.5 L of water and apply thoroughly wet to the bottom 2.5 m of the trunk. Wet the trunk thoroughly but do not spray to run off. Care should be taken to apply the spray right to the base of the root flare.

GARDENS: When damage from cutworms is evident, apply spray to give uniform coverage. Used on broccoli, brussels sprouts, cabbage, chinese cabbage, cauliflower, carrots, celery, cucumber, green pepper, onion, potato, strawberries, sweet corn and tomato. (See table below for days to wait between application and harvest). For carrots and onions use 120 mL per 10 L of water and apply 3 L per 100 m of row. For strawberries use 120 mL in 10 L of water per 100 m2 and apply once between June 1 and June 15. For control of potato flea beetle and tarnished plant bug on potato, apply 80 mL in 10 L of water per 100 m2. Apply weekly as needed. Do not make more than 9 applications during one season. For the other crops listed, use 60 mL per 10 L of water and apply 3 L per 100 m2. Do not use on green bunching onions. Make no more than 1

application per season.

Days to wait between application and harvest:

Broccoli	32 Days	Cucumber	60	Days
Brussels Sprouts	32 Days	Green Peppers	40	Days
Cabbage	32 Days	Onion	60	Days
Carrot	60 Days	Potato	7	Days
Chinese Cabbage	32 Days	Strawberry	20	Days
Cauliflower	32 Days	Sweet Corn	70	Days
Celery	70 Days	Tomato	70	Days

PRECAUTIONS: KEEP OUT OF REACH OF CHILDREN. May be fatal if swallowed. May be absorbed through the skin. May be injurious to eyes and skin. Wash hands and exposed skin after spraying, also any contaminated clothing. Avoid breathing spray mist. Do not permit pets to contact sprayed areas until dry. Do not contaminate food, feed or water; product is poisonous to fish. Re-close container tightly after use and store away from heat.

FIRST AID: Product contains petroleum distillates. If swallowed DO NOT induce vomiting but rush patient to hospital, taking this labelled container with you. If in eyes, flush with plenty of water for 15 minutes, and get medical aid. If on skin wash thoroughly with soap and water.

TOXICOLOGICAL INFORMATION: Chlorpyrifos is a cholinesterase inhibitor; atropine by injection only is antidotal.

DISPOSAL: Dispose of empty container with household garbage. Do not re-use container.

(R) Dursban is a registered trademark of DowElanco

This label transcript service is offered by the Pest Management Regulatory Agency to provide efficient searching for label information. This service and this information do not replace the official hard-copy label. The PMRA does not provide any guarantee or assurance that the information obtained through this service is accurate, current or correct, and is therefore not liable for any loss resulting, directly or indirectly, from reliance upon this service.