

2000/E2001

(( LABEL ))

**READ THE LABEL AND BOOKLET BEFORE USING**

RECOMMENDED AND SOLD IN WESTERN CANADA BY:  
UGG

**United Grain Growers**  
Meeting Farmers' Business Needs

**WEEDONE ®**  
**MCPA ESTER**

**EMULSIFIABLE LIQUID HERBICIDE**

FOR WHEAT, BARLEY, RYE, OATS, FLAX  
PASTURES, STUBBLE LAND, ROADSIDES,  
UNCROPPED LAND AND CONTROL OF WOODY GROWTH

**AGRICULTURAL**

GUARANTEE: MCPA 500 g/L (present as mixed butyl ester)

**CAUTION                      POISON**

REGISTRATION NO. 07825 PEST CONTROL PRODUCTS ACT

**Nufarm**

Nufarm UK Limited  
Crabtree Manorway North  
Belvedere Kent DA17 6BQ England

Canadian Agent:  
Rhone-Poulenc Canada Inc.  
2000 Argentia Road, Plaza 3  
Mississauga, Ontario L5N 1V9  
(905) 821-4450

IN CASE OF SPILLS, POISONING OR FIRE,  
TELEPHONE EMERGENCY RESPONSE NUMBER  
1-800-334-7577 (24 HOURS A DAY)

Weedone® is a registered trademark of Rhone-Poulenc Nederland B.V. - authorized user Rhone-Poulenc Canada Inc.

Rhone-Poulenc Canada Inc. TM is a trademark of Nufarm UK Limited and is used in Canada by Rhone-Poulenc Canada Inc. As exclusive distributor of Nufarm UK Limited.

**PRECAUTIONS:**

KEEP OUT OF REACH OF CHILDREN.

Avoid contact with skin, eyes and clothing.

Wash concentrate from skin or eyes immediately.

Do not inhale fumes.

Avoid breathing spray mist.

After use wash hands and other exposed skin.

Remove and wash contaminated clothing before reuse.

Avoid spray drift.

Store the container tightly closed and away from seeds, feeds, fertilizer, plants and foodstuffs.

**FIRST AID:**

This product contains petroleum distillates.

If swallowed DO NOT induce vomiting. Rush patient to nearest hospital taking this label with you. Never give anything by mouth to an unconscious person.

In case of skin contact, remove contaminated clothing and wash skin thoroughly with soap and water.

In case of eye contact, flush with plenty of water for 15 minutes and get medical attention.

If inhaled, remove to fresh air and get medical attention.

**STORAGE:**

Insecticides and fungicides should be segregated from herbicides so as to prevent the possibility of cross-contamination. This product may be stored at any temperature. SHAKE WELL BEFORE USING.

**DISPOSAL:**

1. Triple rinse the emptied container and add the rinsings to the spray mixture in the tank.
2. Follow provincial instructions for any required additional cleaning of the container prior to its disposal.
3. Make the empty container unsuitable for further use.
4. Dispose of the container in accordance with provincial requirements.
5. For information on the disposal of unused, unwanted product and the cleanup of spills contact the Provincial Regulatory Agency or the Manufacturer.

**NOTICE TO BUYER:** Seller's guarantee shall be limited to the terms set out on the label and subject thereto, the buyer assumes the risk to persons or property arising from the use or handling of this product and accepts the product on that condition.

**NOTICE TO USER:** This control product is to be used only in accordance with the directions on the label. It is an offense under the PEST CONTROL PRODUCTS ACT to use a control product under unsafe conditions.

RECOMMENDED AND SOLD IN WESTERN CANADA BY:

**United Grain Growers (UGG)**

Meetings Farmers' Business Needs

**WEEDONE**

**MCPA ESTER**

**EMULSIFIABLE LIQUID HERBICIDE**

FOR WHEAT, BARLEY, RYE, OATS, FLAX, PASTURES, STUBBLE  
LAND,  
ROADSIDES, UNCROPPED LAND AND CONTROL OF WOODY GROWTH

**AGRICULTURAL**

GUARANTEE: MCPA 500 g/L (present as mixed butyl  
ester)

**CAUTION**

**POISON**

REGISTRATION NO. 7825 PEST CONTROL PRODUCTS ACT

READ THE LABEL AND BOOKLET BEFORE USING

WEDONE® is a registered trademark of Rhone-Poulenc Nederland B.V. - authorized use Rhone-Poulenc Canada Inc. TM is a trademark of Nufarm UK Limited and is used in Canada by Rhone-Poulenc Canada Inc. as exclusive distributor of Nufarm UK Limited.

06/95

**Nufarm**

Nufarm UK Limited  
Crabtree Manorway North

Belvedere Kent DA17 6BQ  
England

**RHONE-POULENC**

Canadian Agent:  
Rhone-Poulenc Canada  
Inc.

2000 Argentia Road, Plaza 3  
Mississauga, Ontario L5N 1V9  
(905) 821-4450

IN CASE OF SPILLS, POISONING OR FIRE, TELEPHONE EMERGENCY  
RESPONSE NUMBER 1-800-334-7577 (24 HOURS A DAY)

RECOMMENDED AND SOLD IN WESTERN CANADA BY:

**United Grain Growers (UGG)**

Meeting Farmers' Business Needs

**WEEDONE®  
MCPA ESTER**

NET

CONTENTS

**10 LITRES**

((BOOKLET))

READ THE LABEL AND BOOKLET BEFORE USING

RECOMMENDED AND SOLD IN WESTERN CANADA BY:

United Grain Growers (UGG)  
Meeting Farmer's Business Needs

**WEEDONE** ®  
**MCPA ESTER**

**EMULSIFIABLE LIQUID HERBICIDE**

FOR WHEAT, BARLEY, RYE, OATS, FLAX  
PASTURES, STUBBLE LAND, ROADSIDES,  
UNCROPPED LAND AND CONTROL OF WOODY GROWTH

**AGRICULTURAL**

GUARANTEE: MCPA 500 g/L  
(present as mixed butyl esters)

**CAUTION                      POISON**

REGISTRATION NO. 7825

PEST CONTROL PRODUCTS ACT

Nufarm  
Nufarm UK Limited  
Crabtree Manorway North  
Belvedere Kent DA17 6BQ  
England

Canadian Agent:  
Rhone-Poulenc Canada Inc.  
2000 Argentinia Road, Plaza 3  
Mississauga, Ontario L5N 1V9  
(905) 821-4450

IN CASE OF SPILLS, POISONING OR FIRE,  
TELEPHONE EMERGENCY RESPONSE NUMBER  
1-800-334-7577 (24 HOURS A DAY)

Weedone(R) is a registered trademark of Rhone-Poulenc  
Nederland B.V. - authorized user Rhone-Poulenc Canada Inc.

TM is a trademark of Nufarm UK Limited and is used in Canada by Rhone-Poulenc Canada Inc. as exclusive distributor of Nufarm UK Limited.

06/95

**PRECAUTIONS:**

KEEP OUT OF REACH OF CHILDREN.

Avoid contact with skin, eyes and clothing.

Wash concentrate from skin or eyes immediately.

Do not inhale fumes.

Avoid breathing spray mist.

After use wash hands and other exposed skin.

Remove and wash contaminated clothing before reuse.

Avoid spray drift.

Store the container tightly closed and away from seeds, feeds, fertilizer, plants and foodstuffs.

**FIRST AID:**

This product contains petroleum distillates.

**If swallowed** DO NOT induce vomiting. Rush patient to nearest hospital taking this label with you. Never give anything by mouth to an unconscious person.

**In case of skin contact**, remove contaminated clothing and wash skin thoroughly with soap and water.

**In case of eye contact**, flush with plenty of water for 15 minutes and get medical attention.

**If inhaled**, remove to fresh air and get medical attention.

**STORAGE:**

Insecticides and fungicides should be segregated from herbicides so as to prevent the possibility of cross-contamination. This product may be stored at any temperature. SHAKE WELL BEFORE USING.

**DISPOSAL:**

1. Triple rinse the emptied container and add the rinsings to the spray mixture in the tank.
2. Follow provincial instructions for any required additional cleaning of the container prior to its disposal.
3. Make the empty container unsuitable for further use.
4. Dispose of the container in accordance with provincial requirements.
5. For information on the disposal of unused, unwanted product and the cleanup of spills contact the Provincial Regulatory Agency or the Manufacturer.

**NOTICE TO BUYER:** Seller's guarantee shall be limited to the terms set out on the label and subject thereto, the buyer assumes the risk to persons or property arising from the use or handling of this product and accepts the product on that condition.

**NOTICE TO USER:** This control product is to be used only in accordance with the directions on the label. It is an offense under the PEST CONTROL PRODUCTS ACT to use a control product under unsafe conditions.

**DIRECTIONS FOR USE:**

**USE PRECAUTIONS:** Avoid spray drift to desirable vegetation **NOT** listed on this label as damage may occur. Avoid contamination of ponds, streams, rivers and other water sources. Do not spray during periods of high winds when spray is likely to drift. Coarse sprays are less likely to drift. Do not spray during periods of high temperatures when crops may be under stress as damage to the crop may occur. Allow 7 days between application of product and grazing or cutting for hay.

**TIMING:** For best results spray when WEEDS ARE YOUNG AND ACTIVELY GROWING. Apply in good growing conditions. Application must be made before the crop canopy shields the weeds.

**SPRAYER:** Clean all spray equipment thoroughly before use.

Select nozzle tips to apply the recommended volume per hectare. Higher water volumes will reduce the risk of crop injury. Calibrate the sprayer and adjust the boom height to ensure uniform coverage. Nozzles must be in good condition and spraying the same volume. Flat fan nozzle tips are recommended.

**MIXING:** Shake this product well before using. Half fill the spray tank with clean water. Add the required amount of product and agitate thoroughly. Fill the tank with clean water and agitate well before use.

**NOTE:** If spray solution has been left standing, agitate thoroughly before use.

**USES:**

**WHEAT, BARLEY, RYE, OATS, (not underseeded with legumes):**

Crops may be sprayed from the full 2-leaf to the early flag leaf (shot blade) stage. Do not apply after the flag leaf stage. Treatment at the 3 to 4-leaf stage usually combines maximum crop tolerance with maximum weed susceptibility. The recommended safe crop rate for wheat, barley and rye is 1.25 litres per hectare. The recommended safe crop rate for oats is 1.1 litres per hectare. Higher rates may be needed to control certain weed species. Possible crop injury from these higher rates is usually offset by the increased yield of the cleaner crop. Use 50 or more litres of water per hectare. Spraying with higher volumes of water (100 to 150 litres per hectare) may reduce the risk of crop injury.

**FALL RYE, WINTER WHEAT:** Treat in the spring between full tillering and the flag leaf (shot blade) stage. DO NOT APPLY DURING AND AFTER THE FLAG LEAF STAGE. Do not apply in the fall. The maximum safe crop rate is 1.25 litres per hectare. Higher rates may be needed to control certain weed species. Possible crop injury from these higher rates is usually offset by the increased yield of the cleaner crop. Use 50 or more litres of water per hectare. Spraying with higher volumes of water (100 to 150 litres per hectare) may reduce the risk of crop injury.

**FLAX:** Apply when the flax is 5 cm tall to just before the bud stage. DO NOT treat after the early bud stage. Due to flax's less competitive nature, weeds should be sprayed in as early a leaf stage as possible, up to the recommended leaf stage. The recommended safe crop rate is 0.85 litres per hectare. Some



crop injury and delay in maturity will occur if flax is treated with more than 0.85 litres of MCPA Ester per hectare. Possible crop injury from these higher rates is usually offset by the increased yield of the cleaner crop. Spraying in the evening and/or increased volumes of water (100 or more litres per hectare) may reduce the risk of flax injury.

**STUBBLE LAND, PASTURES (without legumes), ROADSIDES, UNCROPPED**

**LAND:** Apply at the time of rapid growth (usually May, June and/or September). Spray thoroughly using 1.7 to 3.4 litres of product per hectare in 100 to 300 litres of water. (Use the higher rate for perennial weeds; a second treatment may be required.) Cultivating grain stubble and spraying after regrowth may aid in control. Do not apply on freshly seeded grasses and pasture until established. There is a possibility that plants poisonous to livestock will appear more palatable and could be grazed more readily after MCPA treatment (eg: tansy ragwort, water hemlock, poison hemlock, death camas). Suitable precautions should be taken when such plants are present. Livestock should not be allowed to graze the treated area until poisonous plants have been removed, are dead or have resumed a normal growth habit. Under conditions of good growth and adequate soil moisture, sprays can be applied up to 2 weeks before normal frost time.

**WOODY GROWTH (non-grazing):** Apply at the time of rapid growth (usually May, June and/or September). Apply 2.2 to 4.4 litres of product per hectare in 100 or more litres of water per hectare. Use the lower rate for species that are more susceptible such as chokecherry, western snowberry and willow.

Spray brush species up to 3 metres tall after foliage is well developed. Applications soon after leaves are fully open usually give the best results. Apply sufficient spray volume to thoroughly wet all plant parts to the ground line. Brush or trees above 3 metres tall should be cut close to the ground and sprayed when regrowth is over 0.6 metres tall. With good growing conditions and adequate soil moisture, sprays may be applied up to 2 to 3 weeks before normal frost time.

To reduce potential drift hazard in non-crop locations LO-Drift® may be added. Follow all directions on the LO-Drift label for the correct mixing sequence.

**WEEDS CONTROLLED:**

**SUSCEPTIBLE:** Easy-to-control weeds in the seedling stage (2 to

4 leaf) and growing rapidly. Use 0.8 to 1.1 litres per hectare. Use the higher rate for larger weeds, dry or cold weather, or heavy infestations.

Bluebur, burdock (before the 4-leaf stage), cocklebur, common plantain, flixweed, lamb's-quarters, mustards (except dog and tansy), prickly lettuce, ragweeds, Russian pigweed, shepherd's-purse, stinkweed (field pennycress), wild radish, wild sunflower.

**LESS SUSCEPTIBLE:** Harder-to-control weeds in the seedling stage (2 to 4 leaf) and growing rapidly. Use 1.25 to 2.25 litres per hectare. Use the higher rate for larger weeds, dry or cold weather, or heavy infestations.

Weeds listed under susceptible plus curled dock, dog mustard, hairy galinsoga, oak-leaved goosefoot, hempnettle (before the 4-leaf stage), kochia, peppergrass, purslane,, redroot pigweed, smartweed, tansy mustard.

**TOP GROWTH CONTROL ONLY:** Use 1.25 to 2.25 litres per hectare. Blue lettuce, Canada thistle, dandelion, field bindweed, goat's-beard, gumweed, field bindweed, hedge bindweed, hoary cress, leafy spurge, sow thistle, Tartary buckwheat, wild buckwheat.

**CLEANING OF SPRAYER:** Thoroughly clean the herbicide sprayer before applying any other chemical on crops which may be damaged (see USE PRECAUTIONS). Wash the outside of the sprayer and drain the tank completely. Remove and clean filters, screens and nozzle tips separately. Fill the sprayer tank with clean water. With the agitator running, flush out the lines and boom, then drain. Fill the sprayer with clean water and for each 100L add 1L household ammonia. Operate the pump and agitator for 15 minutes. If possible, let the solution remain in the tank and hoses overnight; recirculate and flush out the lines and boom, then drain. Rinse out twice with clean water, recirculating and draining each time. DO NOT contaminate irrigation or drinking water when cleaning the sprayer.

LO-DRIFT ® is a registered trademark of Rhone-Poulenc Nederland B.V. - autorised user Rhone-Poulenc Canada Inc.

\*\*\*\*\*

This label transcript service is offered by the Pest Management Regulatory Agency to provide efficient searching for label information. This service and this information do not replace the official hard-copy label. The PMRA does not provide any guarantee or assurance that the information obtained through this service is accurate, current or correct, and is therefore not liable for any loss resulting, directly or indirectly, from reliance upon this service.

+))