

23-MAY-1997

((BROCHURE))

INTER-AG
Division of AgrEvo

RINSE For Dollars
RINSE For Safety
RINSE For Disposal

AGRICULTURAL

WARNING POISON

MCPA
Ester 500

GUARANTEE: MCPA 500 g/L
(Present as Ester)

REGISTRATION NO: 21486
PEST CONTROL PRODUCTS ACT

10 LITRES

READ LABEL AND ENCLOSED BROCHURE BEFORE USING

AgrEvo Canada Inc.
295 Henderson Drive
Regina, SK S4N 6C2
Telephone : 1-800-667-5959

PT#LA-M 15 02/97 Printed in USA

LABEL INDEX MCPA ESTER 500	
Wheat, Barley, Rye GO TO Section 1	Filling Instructions GO TO Section 5
Oats and Mixed Grain GO TO Section 2	Precautions GO TO Section 6
Flax GO TO Section 3	First Aid GO TO Section 7
Tank Mixes GO TO Section 4	Rinsing and Disposal GO TO Section 8

SECTION 1	
CROP:	WHEAT, BARLEY, RYE
TREATMENT STAGE:	Treat from the 4th leaf to just before the flag leaf (shot blade) stage. NOTE: Do not treat cereals underseeded to legumes.
SUSCEPTIBLE WEEDS:	Burdock, Cocklebur, Mustards (except Dog Mustard), Ragweed, Flixweeds*, Kochia, Lamb's-quarters, Russian Pigweed, Shepherd's Purse, Stinkweed. * Flixweed susceptible only at seedling stage. Use rates for harder to kill weeds to kill fall germinated flixweed in the spring.
RATES TO USE ON SUSCEPTIBLE WEEDS:	
Small seedlings, good growing conditions:	700 mL/ha
Weeds in bud, dry or cool weather or heavy infestation:	1.1 L/ha
HARDER TO KILL WEEDS:	Annual Sow Thistle, Bluebur, Blue Lettuce*, Canada Thistle*, Peppergrass, Curled Dock, Dog Mustard, Goosefoot, Hempnettle (Suppression only), Redroot Pigweed, Russian Thistle*, Perennial Sow Thistle*, Leafy Spurge*. * Top growth control only.
RATES TO USE ON HARDER TO KILL WEEDS:	
Small seedlings growing rapidly, good growing conditions:	1.25 L/ha
Weeds in bud, dry or cool conditions or heavy infestation:	1.75 L/ha NOTE: Rates over 1.1 L/ha may cause crop injury.
GO TO SECTION 5	
SECTION 2	

CROP:	OATS AND MIXED GRAIN
TREATMENT STAGE:	Spray from 1 - 4 leaf stage.
SUSCEPTIBLE WEEDS:	Burdock, Cocklebur, Mustards (except Dog Mustard), Ragweed, Flixweeds*, Kochia, Lamb's-quarters, Russian Pigweed, Shepherd's Purse, Stinkweed. * Flixweed susceptible only at seedling stage. Use rates for harder to kill weeds to kill fall germinated flixweed in the spring.
RATES TO USE ON SUSCEPTIBLE WEEDS:	
Small seedlings, good growing conditions:	700 mL/ha
Weeds in bud, dry or cool weather or heavy infestation:	1.1 L/ha
HARDER TO KILL WEEDS:	Annual Sow Thistle, Bluebur, Blue Lettuce*, Canada Thistle*, Peppergrass, Curled Dock, Dog Mustard, Goosefoot, Hempnettle (Suppression only), Redroot Pigweed, Russian Thistle*, Perennial Sow Thistle*, Leafy Spurge*. * Top growth control only.
RATES TO USE ON HARDER TO KILL WEEDS:	
Small seedlings growing rapidly, good growing conditions:	1.25 L/ha
Weeds in bud, dry or cool weather or heavy infestation:	1.75 L/ha NOTE: Rates over 1.1 L/ha may cause crop injury.
GO TO SECTION 5	
SECTION 3	
CROP:	FLAX
TREATMENT STAGE:	Spray after plant reaches 5 cm in height and before the bud stage.

SUSCEPTIBLE WEEDS:	Burdock, Cocklebur, Mustards (except Dog Mustard), Ragweed, Flixweed*, Kochia, Lamb's-quarters, Russian Pigweed, Shepherd's Purse, Stinkweed. * Flixweed susceptible only at seedling stage. Use rates for harder to kill weeds to kill fall germinated flixweed in the spring.
RATES TO USE ON SUSCEPTIBLE WEEDS:	
Small seedlings, good growing conditions:	700 mL/ha
Weeds in bud, dry or cool weather or heavy infestation:	1.1 L/ha NOTE: Rates over 850 L/ha may cause crop injury.
GO TO SECTION 5	
SECTION 4	
TANK MIXES with MCPA ESTER 500	Use tank mixes only when they are registered and recommended. Read and follow all label directions on both products used in a tank mix.
TANK MIXES:	MCPA ESTER 500 + Avenge(R) 200-C
CROP:	BARLEY and the following varieties of wheat -NEEPAWA, GLENLEA, CHESTER, SELKIRK, MACOUN, CANUCK, BENITO, COLUMBUS, FIELDER, LEADER, KATEPWA AND HY320.
TREATMENT STAGE:	Apply at the 3 - 5 leaf stage of wild oats.
SUSCEPTIBLE WEEDS:	Wild Oats and Broadleaf weeds - susceptible to MCPA
RATES TO USE:	MCPA ESTER 500 1.1 L/ha + Avenge® 200-C + 3.5 - 4.25 L/ha Apply this mixture in 100 Litres of water per hectare.
GO TO SECTION 5	
SECTION 5	

FILLING SPRAY TANK	<ol style="list-style-type: none">1. Fill tank 1/2 full with water.2. Add MCPA Ester 500. When Avenge is used, add after MCPA Ester 500.3. Complete filling of tank with water.4. Use 60 - 240 litres of water per hectare unless otherwise specified.5. Do not exceed 275 kPa pressure.
GO TO SECTION 6	
SECTION 6	

PRECAUTIONS:

1. Keep out of the reach of children.
2. Avoid contact with eyes, skin and clothing. May cause skin irritation.
3. Do not breathe spray mist or vapors. Avoid spray or vapor drifts onto susceptible plants such as grapes, tomatoes, tobacco, sugar beets, sunflowers, beans, turnips, cauliflower, cabbage, alfalfa, clover, peas, potatoes, ornamental trees, shrubs or flowers, fruit trees, lawns, most vegetable crops or other desirable vegetation.
4. Do not spray when there is any danger of wind or on exceptionally hot days over 27°C.
5. Drift from spray may be reduced by using high volume sprays under low pressure, coarse sprays, and drop nozzles. Use special care in airplane application where danger from drift is greater.
6. For aerial application, exercise special care:
 - 1) Do not spray in winds exceeding 8 km per hour.
 - 2) Do not spray in dead calm near sensitive plants. A cloud of suspended droplets may drift onto sensitive plants when the wind comes up.
 - 3) Spray only when the wind is blowing away from a sensitive crop, shelter belt or garden.
7. Do not contaminate any body of water. Avoid contamination of foods.
8. Do not use in a greenhouse.
9. Keep in original container during storage.
10. Do not store near or in same room as seeds, feeds, fertilizers or pesticides.
11. Keep away from fire or open flame and other sources of heat.
12. Clean spray equipment thoroughly after use. Do not use spray equipment for other purposes. Do not re-use empty containers.
13. Allow 7 days between application of product and grazing/cutting for hay.

GO TO SECTION 7

SECTION 7

FIRST AID:

1. In case of poisoning, call a physician or contact a poison control centre immediately.
2. If on skin, remove contaminated clothing and wash skin thoroughly with soap and water.
3. If in eyes, flush with water immediately and continue for 5-10 minutes. Get medical attention or contact a poison control centre.
4. If swallowed, do not induce vomiting. Take patient to nearest hospital or doctor's office, taking the pesticide label with you, or contact a poison control centre.

GO TO SECTION 8

SECTION 8

DISPOSAL INFORMATION:

RINSE For Dollars
RINSE For Safety
RINSE For Disposal

1. Triple rinse the emptied container thoroughly and add the rinsings to the spray mixture in the tank.
2. Follow provincial instructions for any required additional cleaning of the container prior to its disposal.
3. Make the empty container unsuitable for further use.
4. Dispose of the container in accordance with provincial requirements.
5. For information on the disposal of unused, unwanted product and the cleanup of spills, contact the Provincial Regulatory Agency or the Manufacturer.

NOTICE TO USER:

This control product is to be used only in accordance with the directions on this label. It is an offense under the Pest Control Products Act to use a control product under unsafe conditions.

NOTICE TO BUYER:

Seller's guarantee shall be limited to the terms set out on the label, and subject thereto the buyer assumes the risk to persons or property arising from the use or handling of this product and accepts the product on that condition.

AgrEvo, and INTER-AG. are registered trademarks of HOECHST SCHERING
AGREVO GmbH, Germany.
AVENGE is a registered trademark of Cyanamid Canada Inc.

((LABEL))

INTER-AG
Division of AgrEvo

RINSE For Dollars
RINSE For Safety
RINSE For Disposal

AGRICULTURAL

WARNING POISON

MCPA
Ester 500

GUARANTEE: MCPA 500 g/L
(Present as Ester)

REGISTRATION NO. 21486
PEST CONTROL PRODUCTS ACT

10 LITRES

READ LABEL AND ATTACHED DIRECTIONS FOR USE BEFORE USING

AgrEvo Canada Inc.
295 Henderson Drive
Regina, SK S4N 6C2
Telephone : 1-800-667-5959

PRECAUTIONS

1. Keep out of the reach of children.
2. Avoid contact with eyes, skin and clothing. May cause skin irritation.
3. Do not breathe spray mist or vapours. Avoid spray or vapour drifts onto susceptible plants such as grapes, tomatoes, tobacco,

sugar beets, sunflowers, beans, turnips, cauliflower, cabbage, alfalfa, clover, peas, potatoes, ornamental trees, shrubs or flowers, fruit trees, lawns, most vegetable crops or other desirable vegetation.

4. Do not spray when there is any danger of wind or on exceptionally hot days over 27°C.
5. Drift from spray may be reduced by using high volume sprays under low pressure, coarse sprays, and drop nozzles. Use special care in airplane application where danger from drift is greater.
6. For aerial application, exercise special care:
 - 1) Do not spray in winds exceeding 8 km per hour.
 - 2) Do not spray in dead calm near sensitive plants. A cloud of suspended droplets may drift onto sensitive plants when the wind comes up.
 - 3) Spray only when the wind is blowing away from a sensitive crop, shelter belt or garden.
7. Do not contaminate any body of water. Avoid contamination of foods.
8. Do not use in a greenhouse.
9. Keep in original container during storage.
10. Do not store near or in same room as seeds, feeds, fertilizers or pesticides.
11. Keep away from fire or open flame and other sources of heat.
12. Clean spray equipment thoroughly after use. Do not use spray equipment for other purposes. Do not re-use empty containers.
13. Allow 7 days between application of product and grazing/cutting for hay.

DISPOSAL INFORMATION

1. Triple rinse the emptied container thoroughly and add the rinsings to the spray mixture in the tank.
2. Follow provincial instructions for any required additional cleaning of the container prior to its disposal.
3. Make the empty container unsuitable for further use.
4. Dispose of the container in accordance with provincial requirements.
5. For information on the disposal of unused, unwanted product and the cleanup of spills, contact the Provincial Regulatory Agency or the Manufacturer.

FIRST AID:

1. In case of poisoning, call a physician or contact a poison control centre immediately.
2. If on skin, remove contaminated clothing and wash skin thoroughly with soap and water.
3. If in eyes, flush with water immediately and continue for 5-10 minutes. Get medical attention, or contact a poison control centre.
4. If swallowed, do not induce vomiting then take patient to nearest

hospital or doctor's office, taking the pesticide label with you,
or contact a poison control centre.

NOTICE TO USER:

This control product is to be used only in accordance with the
directions on this label. It is an offence under the Pest Control
Products Act to use a control product under unsafe conditions.

23-MAY-1997

**IMPORTANT: LABEL CORRECTION
ATTENTION: FLAX GROWERS**

THE LABEL ATTACHED TO THE JUGS, INSIDE THIS CARTON, CONTAINS INCORRECT APPLICATION RATES FOR FLAX. IF USING THIS PRODUCT ON FLAX, USE THE RATES INDICATED BELOW.

SECTION 3	
Crop:	Flax
TREATMENT STAGE:	Spray after plant reaches 5 cm in height and before the bud stage.
Susceptible Weeds:	Burdock, Cocklebur, Mustards (except Dog Mustard), Ragweed, Flixweed*, Kochia, Lamb's-quarters, Russian Pigweed, Shepherd's Purse, Stinkweed. <i>*Flixweed susceptible only at seedling stage. Use rates for harder to kill weeds to kill fall germinated flixweed in the spring.</i>
Rates to Use on Susceptible Weeds:	
Small seedlings, good growing conditions:	700 mL/ha
Weeds in bud, dry or cool weather or heavy infestation:	1.1 L/ha NOTE: Rates over 850 mL/ha may cause crop injury.

97.05.23

**IMPORTANT: LABEL CORRECTION
ATTENTION: FLAX GROWERS**

THE LABEL ATTACHED TO THE JUGS, INSIDE THIS CARTON, CONTAINS INCORRECT APPLICATION RATES FOR FLAX. IF USING THIS PRODUCT ON FLAX, USE THE

RATES INDICATED BELOW.

SECTION 3	
Crop:	Flax
TREATMENT STAGE:	Spray after plant reaches 5 cm in height and before the bud stage.
Susceptible Weeds:	Burdock, Cocklebur, Mustards (except Dog Mustard), Ragweed, Flixweed*, Kochia, Lamb's-quarters, Russian Pigweed, Shepherd's Purse, Stinkweed. <i>*Flixweed susceptible only at seedling stage. Use rates for harder to kill weeds to kill fall germinated flixweed in the spring.</i>
Rates to Use on Susceptible Weeds:	
Small seedlings, good growing conditions:	700 mL/ha
Weeds in bud, dry or cool weather or heavy infestation:	1.1 L/ha NOTE: Rates over 850 mL/ha may cause crop injury.

23-MAY-1997

GUARANTEE: MCPA 500 g/L (Present as Ester)
REGISTRATION NO. 21486
PEST CONTROL PRODUCTS ACT

AgrEvo Canada Inc.
295 Henderson Drive, Regina, SK, S4N 6C2
Telephone: 1-800-667-5959

**MCPA
ESTER 500
10 LITRES**

WARNING

POISON

SUPPLEMENTAL LABEL

READ LABEL AND BROCHURE PRIOR TO USING

AGRICULTURAL

IMPORTANT: LABEL CORRECTION

ATTENTION: FLAX GROWERS

The Label attached to the jugs, inside this carton, contains incorrect application rates for Flax. If using this product on Flax, use the rates indicated below.

SECTION 3	
Crop:	FLAX
Treatment Stage:	Spray after plant reaches 5 cm in height and before the bud stage.
Susceptible Weeds:	Burdock, Cocklebur, Mustards (except Dog Mustard), Ragweed, Flixweed*, Kochia, Lamb's-quarters, Russian Pigweed, Shepherd's Purse, Stinkweed. <i>*Flixweed susceptible only at seedling stage. Use rates for harder to kill weeds to kill Fall germinated Flixweed in the Spring.</i>
Rates to Use on Susceptible Weeds:	
Small seedlings, good growing conditions:	700 mL/ha
Weeds in bud, dry or cool weather or heavy infestation:	1.1 L/ha NOTE: Rates over 850 mL/ha may cause crop injury.

PRECAUTIONS

1. Keep out of the reach of children.
2. Avoid contact with eyes, skin and clothing. May cause skin irritation.
3. Do not breathe spray mist or vapours. Avoid spray or vapour drifts onto susceptible plants such as grapes, tomatoes, tobacco, sugar beets, sunflowers, beans, turnips, cauliflower, cabbage, alfalfa, clover, peas, potatoes, ornamental trees, shrubs or flowers, fruit trees, lawns, most vegetable crops or other desirable vegetation.
4. Do not spray when there is any danger of wind or on exceptionally

hot days over 27°C.

5. Drift from spray may be reduced by using high volume sprays under low pressure, coarse sprays, and drop nozzles. Use special care in airplane application where danger from drift is greater.
6. For aerial application, exercise special care: **1)** Do not spray in winds exceeding 8 km per hour. **2)** Do not spray in dead calm near sensitive plants. A cloud of suspended droplets may drift onto sensitive plants when the wind comes up. **3)** Spray only when the wind is blowing away from a sensitive crop, shelter belt or garden.
7. Do not contaminate any body of water. Avoid contamination of foods.
8. Do not use in a greenhouse.
9. Keep in original container during storage.
10. Do not store near or in same room as seeds, feeds, fertilizers or pesticides.
11. Keep away from fire or open flame and other sources of heat.
12. Clean spray equipment thoroughly after use. Do not use spray equipment for other purposes. Do not re-use empty containers.
13. Allow 7 days between application of product and grazing/cutting for hay.

DISPOSAL INFORMATION

1. Triple rinse the emptied container thoroughly and add the rinsings to the spray mixture in the tank.
2. Follow provincial instructions for any required additional cleaning of the container prior to its disposal.
3. Make the empty container unsuitable for further use.
4. Dispose of the container in accordance with provincial requirements.
5. For information on the disposal of unused, unwanted product and the cleanup of spills, contact the Provincial Regulatory Agency or the Manufacturer.

FIRST AID:

1. In case of poisoning, call a physician or contact a poison control centre immediately.
2. If on skin, remove contaminated clothing and wash skin thoroughly with soap and water.
3. If in eyes, flush with water immediately and continue for 5-10 minutes. Get medical attention, or contact a poison control centre.
4. If swallowed, induce vomiting then take patient to nearest hospital or doctor's office. Take the pesticide label with you, or contact a poison control centre.

NOTICE TO USER: This control product is to be used only in accordance with the directions on this label. It is an offence under the Pest Control Products Act to use a control product under unsafe conditions.

This label transcript service is offered by the Pest Management Regulatory Agency to provide efficient searching for label information. This service and this information do not replace the official hard-copy label. The PMRA does not provide any guarantee or assurance that the information obtained through this service is accurate, current or correct, and is therefore not liable for any loss resulting, directly or indirectly, from reliance upon this service.

+))