

94.12.02

DOMESTIC

REG. NO. 23584
P.C.P. ACT

CROAK TM
PELLETED BAIT
WITH
cov-R-tox TM
Encapsulated Warfarin

KILLS
Rats & Mice

For Control of Rats and Mice
Indoors and Outdoors

READ THE LABEL BEFORE USING

GUARANTEE: Warfarin.....0.025%

NET: 453.6 g

HACCO, Inc.
P.O. Box 7190
Madison, WI 53707

DIRECTIONS: NORWAY AND ROOF RATS: Use approximately 100-450 grams per bait station in locations frequented by rats or where rats have been know to feed. Baits must be protected from pets, poultry, livestock, etc., by bait boxes or other suitable means. Baits should be examined regularly and replenished before completely eaten. Replace bait if it molds or decomposes.

HOUSE MICE: For controlling mice, use 40-60 grams per bait station and place 3-4 meters apart. Follow the same general directions as for rats, except that the amount of bait placements should be smaller and more placements should be made. Mice are more difficult to control than rats and complete control may take a longer period of baiting.

IMPORTANT: Baiting should be continued as long as evidence of feeding is observed. Assure an uninterrupted supply of fresh bait for a period of not less than 12 days. Where a continuous source of infestation is present from nearby dumps or fields,

permanent stations should be used and bait placed in them as often as necessary.

PRECAUTIONS: KEEP OUT OF REACH OF CHILDREN AND PETS

FIRST AID: CROAK TM Pelleted Bait contains as its active ingredient an anticoagulant chemical which, if taken accidentally by human, domestic animals or pets, may reduce the clotting ability of the blood and internal hemorrhage may result. In all case baits are accidentally eaten, give patient 1-2 glasses of water and cause vomiting by inserting of a finger down the throat, repeat until vomit fluid is clear. Call a Physician or Poison Control Centre immediately.

TOXICOLOGY: When a human has been know to have accidentally ingested CROAK TM Pelleted Bait, blood transfusions combined with intravenous injections and oral doses of Vitamin K1 are indicated as in the case of hemorrhage caused by overdoses of Dicumarol. Vitamin K1 in the form of intravenous intramuscular or subcutaneous injections, or by oral ingestion are suggested remedial treatments for anticoagulant poisoning. The severity of the case measured by establishing prolonged prothrombin times (P.T.) will determine appropriate therapy. Monitoring P.T. will indicate the necessity of repeated treatments.

ENVIRONMENTAL HAZARDS: Keep out of lakes, streams, or ponds.

DISPOSAL: Do not reuse container. Wrap and dispose of with household garbage.

NOTICE TO BUYER: Seller's guarantee shall be limited to the terms set out in the label and subject thereto, the buyer assumes the risk to persons or property arising from the use or handling of this product and accepts the product on that condition.

This label transcript service is offered by the Pest Management Regulatory Agency to provide efficient searching for label information. This service and this information do not replace the official hard-copy label. The PMRA does not provide any guarantee or assurance that the information obtained through this service is accurate, current or correct, and is therefore not liable for any loss resulting, directly or indirectly, from reliance upon this service.

+))