

07-FEB-2007

CARTON

**HOME
WARFARIN
RAT AND MOUSE KILLER MEAL**

RODENTICIDE

DOMESTIC

REG. NO. 15739 P.C.P. ACT

GUARANTEE: WARFARIN 0.025%

Contains malathion at 0.0095% and calcium propionate at 0.3% as preservatives.

**READ THE LABEL BEFORE USING
KEEP OUT OF THE REACH OF CHILDREN**

500 g (1 kg)
2 (4) Ready-to-Use Bait Bags

**HOME HARDWARE STORES LIMITED
34 Henry St. W.
ST. JACOBS, ONTARIO N0B 2N0
Tel: 519-664-2252**

HOME Warfarin contains an anticoagulant chemical that reduces the clotting ability of the blood. After several feedings rodents die from fatal haemorrhage.

DIRECTIONS FOR USE: Place bait feeding stations in locations frequented by rats and mice, near rat holes, runways or burrows; or against walls or in corners where mice are seen.

Place bait in tamper-resistant bait boxes or protected areas inaccessible at the time of placement to children, pets, domestic animals and wildlife.

A shallow container of drinking water encourages consumption of bait.

For Rats: Norway rats (*Rattus norvegicus*) and Roof rats (*Rattus rattus*). Place 250 - 500 g of bait at 4 - 5 m intervals. Maintain a constant supply of bait at stations for at least 10 days.

For Mice: (*Mus musculus*) Place 250 g of bait at 4 - 5 m intervals. An uninterrupted supply of bait should be maintained for at least 15 days.

PRECAUTIONS: KEEP OUT OF REACH OF CHILDREN, PETS AND LIVESTOCK. May be harmful or fatal if swallowed or absorbed through the skin. Do not open pre-measured place packs. Rubber gloves must be worn when disposing of dead rodents, unconsumed bait and empty containers. Avoid contact with eyes, skin or clothing. Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet. Wash skin thoroughly with soap and water after handling. Wash contaminated clothing, separately from other laundry, with soap and hot water before reuse. KEEP AWAY FROM FEED AND FOODSTUFFS. Keep out of lakes, streams or ponds.

USE LIMITATIONS: Bait MUST either be placed in tamper-resistant bait stations or in locations not accessible to children, pets or livestock. DO NOT place bait in areas where there is a possibility of contaminating food or surfaces that come in direct contact with food.

FIRST AID: This product contains an anticoagulant chemical which, if taken accidentally by humans, domestic animals, pets or wildlife may reduce the clotting ability of the blood and haemorrhage may result. **If in eyes,** flush with water for 15 minutes. **If on skin,** wash with soap and water. **If swallowed,** give patient one or two glasses of water and induce vomiting by touching back of the throat with a finger or blunt object; repeat until vomit fluid is clear. Do not give anything by mouth to an unconscious person. For all cases of human ingestion, immediately notify a physician or poison control centre. Take container, label or product name and P.C.P. Registration Number with you when seeking medical attention. If pet or livestock poisoning is suspected, immediately contact a veterinarian.

TOXICOLOGICAL INFORMATION: Vitamin K₁ in the form of intramuscular or subcutaneous injections, or by oral ingestion are suggested remedial treatments for anticoagulant poisoning. The severity of the case measured by establishing prolonged prothrombin times (P.T.) will determine appropriate therapy. Monitoring P.T. will indicate the necessity of repeated treatments.

STORAGE: Store in a cool, dry place away from other chemicals and food or feed. Store product not in use, in original container, in a secure location inaccessible to children and non-target animals.

DISPOSAL: Do not reuse empty container. Dispose of empty container by placing in household garbage. Dispose of dead rodents in garbage or by burying.

NOTICE TO USER: This pest control product is to be used only in accordance with the directions on the label. It is an offence under the Pest Control Products Act to use this product in a way that is inconsistent with the directions on the label. The user assumes the risk to persons or property that arises from any such use of this product.

07-FEB-2007

THROW BAG:

**HOME
WARFARIN
RAT AND MOUSE KILLER MEAL**

RODENTICIDE

DOMESTIC

REG. NO. 15739 P.C.P. ACT

GUARANTEE: WARFARIN 0.025%

Contains malathion at 0.0095% and calcium propionate at 0.3% as preservatives.

**READ THE LABEL BEFORE USING
KEEP OUT OF THE REACH OF CHILDREN**

250 g

**HOME HARDWARE STORES LIMITED
34 Henry St. W.
ST. JACOBS, ONTARIO N0B 2N0
Tel: 519-664-2252**

PRECAUTIONS: KEEP OUT OF REACH OF CHILDREN, PETS AND LIVESTOCK.

May be harmful or fatal if swallowed or absorbed through the skin. Do not open pre-measured place packs. Rubber gloves must be worn when disposing of dead rodents, unconsumed bait and empty containers. Avoid contact with eyes, skin or clothing.

Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.

Wash skin thoroughly with soap and water after handling. Wash contaminated clothing, separately from other laundry, with soap and hot water before reuse. **KEEP AWAY FROM FEED AND FOODSTUFFS.** Keep out of lakes, streams or ponds.

USE LIMITATIONS: Bait **MUST** either be placed in tamper-resistant bait stations or in locations not accessible to children, pets or livestock. **DO NOT** place bait in areas where there is a possibility of contaminating food or surfaces that come in direct contact with food.

FIRST AID: This product contains an anticoagulant chemical which, if taken accidentally by humans, domestic animals, pets or wildlife, may reduce the clotting ability of the blood and haemorrhage may result. **If in eyes,** flush with water for 15 minutes. **If on skin,** wash with soap and water. **If swallowed,** give patient one or two glasses of water and induce vomiting by touching back of the throat with a finger or blunt object; repeat until vomit fluid is clear. Do not give anything by mouth to an unconscious person. For all cases of human ingestion, immediately notify a physician or poison control centre. Take container, label or product name and P.C.P. Registration Number with you when seeking medical attention. If pet or livestock poisoning is suspected, immediately contact a veterinarian.

TOXICOLOGICAL INFORMATION: Vitamin K₁ in the form of intramuscular or subcutaneous injections, or by oral ingestion are suggested remedial treatments for anticoagulant poisoning. The severity of the case measured by establishing prolonged prothrombin times (P.T.) will determine appropriate therapy. Monitoring P.T. will indicate the necessity of repeated treatments.

STORAGE: Store in a cool, dry place away from other chemicals and food or feed. Store product not in use, in original container, in a secure location inaccessible to children and non-target animals.

DISPOSAL: Do not reuse empty container. Dispose of empty container by placing in household garbage. Dispose of dead rodents in garbage or by burying.

This label transcript service is offered by the Pest Management Regulatory Agency to provide efficient searching for label information. This service and this information do not replace the official hard-copy label. The PMRA does not provide any guarantee or assurance that the information obtained through this service is accurate, current or correct, and is therefore not liable for any loss resulting, directly or indirectly, from reliance upon this service.