## GREEN CROSS(R)

## BUG-B-GON (R) ANT AND GRUB KILLER

### DOMESTIC

### INSECTICIDE DUST

Controls
ants, grubs, maggots, cutworms and other
insects in gardens and lawns.

### **GUARANTEE:**

This product contains the DURSBAN\* brand of chlorpyrifos.

\*DURSBAN is a registered trademark of DowElanco Canada Inc.

REG. NO.: 18801 P.C.P. ACT

NET WEIGHT: 500 g

READ THE LABEL BEFORE USING

KEEP OUT OF REACH OF CHILDREN

The Solaris Group
A Unit of Monsanto Canada Inc.
600 - 25 Watline Avenue
Mississauga, Ontario
L4Z 2Z1

### DIRECTIONS:

Remove child resistant cap with a coin or tool. Shake to apply. Reapply cap after use.

FOR USE ON	TO CONTROL	APPLICATION RATE/	TIMING
		AREA OF COVERAGE	

LAWNS:	White grubs,	Apply 400 g per	Make no more than 2		
	European chafer larvae, Japanese beetle larvae	10 sq. m(14 oz per 110 sq. ft.) and water into turf.	applications per season. KEEP OFF LAWN UNTIL DRY.		
	Ants	Apply 200 g per 10 sq. m (7 oz per 110 sq. ft).	Treat ant nests and trails. Make no more than 2 applications per season. KEEP OFF TREATED AREA UNTIL DUST HAS DISSIPATED.		
	Chinch bugs	Apply at a rate of 200 g per 10 sq. m (7 oz per 110 sq. ft) and water into turf.	Apply at first sign of damage. Water thoroughly. Make no more than 2 applications per season. KEEP OFF TREATED AREA UNTIL DRY.		
GARDENS: Broccoli, Brussels, sprouts, cabbage, Chinese cabbage, cauliflower, carrots, celery, cucumber, green pepper, onion, potato, sweet corn and tomato. DO NOT USE ON GREEN BUNCHING ONIONS.	Cutworms, root maggots	Apply dust to soil around plant at a rate of 70 g per 10 meters (2.5 oz per 33 feet) of row if damage is expected.	Chemical control of white grubs and wireworms is usually needed only the year after sod has been turned.		
DAYS TO WAIT BETWEEN APPLICATION AND HARVEST					
Crop: Potato Potato  Broccoli, Brussels sprouts, Cabbage, Cauliflower, Chinese Cabbage Green pepper Carrot, Cucumber, Onion Celery, Sweet corn, Tomato  7  7  7  7  7  7  7  7  7  7  7  7  7					

ORNAMENTALS:	spiders, cutworms, white	or to soil around plants at a rate of 400 g per 10 square meters	Make no more than 2 applications per season.
--------------	--------------------------	--	--

### PRECAUTIONS:

Do not allow dust to contact food, food preparing surfaces, utensils or drinking water. Wash contaminated food surfaces and utensils with soap and water. Do not use in poultry houses or other animal buildings. May cause irritation to skin or eyes; avoid contact. Avoid inhaling dust. Wash thoroughly with soap and water after using. KEEP OUT OF REACH OF CHILDREN.

### WARNING:

This product is toxic to fish and crustaceans. Avoid contamination of ornamental fish ponds and fish bowls. Dispose of the container away from food, feed and water supplies.

#### FIRST AID:

If swallowed, induce vomiting and seek medical attention or contact a Poison Control Centre. Flush contaminated eyes and skin with plenty of water. For eyes get prompt medical attention.

# TOXICOLOGICAL INFORMATION:

Chlorpyrifos is a cholinesterase inhibitor. Treat symptomatically. Atropine by injection only is an antidote.

### DISPOSAL:

Do not reuse container. Wrap and dispose of empty container with household garbage.

### NOTICE TO BUYER:

Seller's guarantee shall be limited to the terms set out on the label and subject thereto, the buyer assumes the risk to persons or property arising from the use or handling of this product and accepts the product on that condition.

Product Use Information: 1-800-668-5669

(R)Registered trademark of Monsanto Canada Inc. licensee

This label transcript service is offered by the Pest Management Regulatory Agency to provide efficient searching for label information. This service and this information do not replace the official hard-copy label. The PMRA does not provide any guarantee or assurance that the information obtained through this service is accurate, current or correct, and is therefore not liable for any loss resulting, directly or indirectly, from reliance upon this service.

+))