

2001/2002

KE

**MARLATE\* 50  
INSECTICIDE**

**WETTABLE POWDER FOR CROPS AND LIVESTOCK**

**AGRICULTURAL**

**READ THE LABEL BEFORE USING**

REGISTRATION NO. 05054 - PEST CONTROL PRODUCTS ACT

**GARANTEE:** Methoxychlor Technical ..... 50%

**NET CONTENTS: Kg**

**PRECAUTIONS:**

KEEP OUT OF REACH OF CHILDREN

Avoid contact with skin, eyes, clothing.  
Avoid breathing spray mist or dust.  
Avoid contamination of feed and food.  
Wash thoroughly after handling, or using.  
Do not contaminate any body of water.  
Do not contaminate mangers, drinking troughs or utensils.

**FIRST AID:** For external contact, wash with soap and water.  
If in eyes, flush with plenty of water. Get medical attention. If swallowed, drink 1 or 2 glasses of water and induce vomiting by touching back of throat with finger, or blunt object. Do not induce vomiting or give anything by mouth to an unconscious person. Get medical attention.

Do not re-use bag - bury when empty.

\*MARLATE KE Registered Trademark in Canada.

Manufactured by

**KINCAID ENTERPRISES, INC.  
P.O. Box 549  
Nitro, West Virginia 25143**

## USA

### DIRECTIONS AGRICULTURAL CROPS

KE Marlate\* 50 Insecticide should be used only in accordance with recommendations on this label, or in separate published KE recommendations available through local dealers.

Use sufficient spray or dust to thoroughly cover the plant. Start applications at first sign of infestation and repeat at 7-14 day intervals or as necessary. Do not apply during bloom of fruit crops.

**SPRAYS** - For fruit, apply 2.5-4 kg "Marlate" 50 per 1000 L water (2.5-4 g. per litre of water); Garden Crops 2-5 kg per ha (asparagus 1.75-4 kg).

**DUSTS** - For small fruits and garden crops except asparagus and tomatoes, make a 6-20% dust (3-10% active) by thoroughly mixing "Marlate" 50 with any suitable diluent, preferably talc or pyrophyllite.

**TREE FRUITS** - Cherry fruit fly, plum curculio.

**Apples, cherries, pears, plums:** preharvest interval - 7 days.

**Peaches, apricots:** 21 days.

Make 2 or 3 applications after petal fall. Follow provincial recommendations for timing.

**SMALL FRUITS** - Currant fruit fly, strawberry weevil.

**Strawberry, currant, gooseberry:** preharvest interval - 14 days.

**GARDEN CROPS** - Asparagus beetle, bean weevil, bean leaf beetle, leafhoppers, Mexican bean beetle, blister beetle, flea beetles, cabbage looper, diamond-back moth, imported cabbage worm, cucumber beetle, squash bug, thrips, pea weevil,

Colorado potato beetle, fall armyworm, tomato fruitworm.

**Carrot\*, cucumber, eggplant\*, kohlrabi, melon, pepper\*, pumpkin, turnip\*, squash, tomato:** preharvest interval - 1 day.

**Asparagus, cabbage:** preharvest interval - 3 days.

**Bean\*, beet\*, cauliflower, onion, pea\*, potato\*, radish\*, rutabaga\*, turnip, carrot:** preharvest interval - 7 days. Do not use on young lima bean plants.

**Broccoli, brussels sprouts, kale, lettuce\*, spinach\*:** preharvest interval - 14 days.

\* Do not feed treated crop or crop refuse to livestock.

**FIELD AND FORAGE CROPS** - Alfalfa weevil (larvae), flea beetles, grasshoppers, leafhoppers, sweet clover weevil, alfalfa blotch leafminer.

**Alfalfa, clover** - apply 2.5-2.75 kg per ha. For armyworm control apply 4.5 to 5.5 kg per ha. Preharvest and grazing interval - 7 days. May be applied during bloom at early morning or late evening when bees are not foraging.

#### **BEEF CATTLE, HORSES, SHEEP, GOATS**

Do not apply within 7 days of slaughter.

**BEEF CATTLE** - Hornflies. Treatments are usually effective for about three weeks. Do not apply to dairy animals.

**Spray or Dip** - Use 0.5-1.0 kg per 100 L water. As a spray, make sure animals are wetted to the skin. Do not dip calves under 1 month of age. Use lower rate on young animals and calves.

**Dust** - Use 10% dust. Apply to the poll, neck and upper sides and rub in lightly to carry powder beneath the hair.

**HORSES** - Fleas, mosquitoes, stable fly.

**Spray** - Use 1.0-2.0 kg per 100 L water. Spray 30 to 60 mL per animal and repeat daily if necessary. Use lower rate on young colts.

**GOATS, SHEEP** - lice, ticks (keds). Do not use on young animals.

**Spray** - Use 1.0-2.0 kg per 100 L water. Spray 30 to 60 mL per animal after shearing and repeat at 2 week intervals.

**Dust** - Use 10% dust after shearing and lightly cover the entire animal at 2 week intervals.

**NOTICE TO USER:** This control product is to be used only in accordance with the directions on this label. It is an offense under the PEST CONTROL PRODUCTS ACT to use a control product under unsafe conditions.

**NOTICE TO BUYER:** Seller' s guarantee shall be limited to the terms set out on the label and subject thereto, the Buyer assumes the risk to persons or property arising from the use of handling of this product and accepts the product on that condition.

Made in U.S.A.

Printed in U.S.A.

\*\*\*\*\*

This label transcript service is offered by the Pest Management Regulatory Agency to provide efficient searching for label information. This service and this information do not replace the official hard-copy label. The PMRA does not provide any guarantee or assurance that the information obtained through this service is accurate, current or correct, and is therefore not liable for any loss resulting, directly or indirectly, from reliance upon this service.

+))