

86.10.03

(( LABEL ))

SANEX

VAPO 20 E.C.  
Emulsifiable Concentrate Insecticide

COMMERCIAL

DANGER POISON

READ THE LABEL BEFORE USING

500 millilitres

REGISTRATION NO. 12539 PEST CONTROL PRODUCTS ACT

GUARANTEE: Dichlorvos.....20%

Manufactured by  
SANEX INC.  
2965 Slough Street  
Mississauga, Ontario  
L4T 1G2

Branches throughout North America

DIRECTIONS

(SEE LEAFLET FOR COMPLETE DIRECTIONS)

PRECAUTIONS: Do not get on skin or eyes. Do not breathe fumes. Wear natural rubber gloves and protective clothing when handling. Wear a mask or respirator offering protection against dichlorvos insecticide. Keep in original container during storage. Do not store near nor contaminate feed or foodstuffs. Do not contaminate streams, lakes, ponds, irrigation water, water used by livestock or water used for domestic purposes. Keep product away from flame or heat.

FIRST AID: In case of skin contact, wash immediately with soap and water. Wash all contaminated clothing before re-use. In case of eye contact, flush eyes with running water for 10 minutes. Product contains petroleum distillate. If accidentally swallowed, do not induce vomiting. Keep victim still and quiet. Give artificial respiration if victim is not breathing. Call a

doctor immediately.

NOTE: This product contains petroleum distillates.

TOXICOLOGY: Symptoms of poisoning include weakness, headache, tightness in chest, blurred vision, non-reactive pin-point pupils, salivations, sweating, nausea, vomiting, diarrhea, and abdominal cramps. If there are signs of parasympathetic stimulation, atropine sulphate should be injected at 10 minute intervals. In doses of 1 to 2 milligrams until complete atropinization has occurred. Morphine is contraindicated. Clear chest by postural drainage. Observe patient continuously for 48 hours. Repeated exposure to cholinesterase inhibitors may, without warning, cause prolonged susceptibility to very small doses of any cholinesterase inhibitor. Allow no further exposure until time for cholinesterase regeneration has been allowed as determined by blood test.

NON WARRANTY: Seller's guarantee shall be limited to the terms set out on the label and subject thereto, the buyer assumes the risk to persons or property arising from the use or handling of this product and accepts the product on that condition.

NOTICE TO USER: This control product is to be used only in accordance with the directions on this label. It is an offence under the Pest Control Products Act to use a control product under unsafe conditions.

(( PAMPHLET ))

SANEX

VAPO 20

Emulsifiable Concentrate Insecticide

COMMERCIAL

DANGER

POISON

READ THE LABEL BEFORE USING

GUARANTEE: Dichlorvos.....20%

NET CONTENTS 568 mL

REGISTRATION NO. 12539 PEST CONTROL PRODUCTS ACT

Manufactured by  
SANEX INC.

2965 Slough Street  
Mississauga, Ontario  
L4T 1G2

Branches throughout North America

#### DIRECTIONS

DAIRY CATTLE, BEEF CATTLE, HORSES: For the control of face flies thoroughly mix 5 mL of Sanex Vapo 20 Emulsifiable Insecticide with 85 mL of water and 85 mL of corn syrup. Apply 5.5 mL of this bait to the forehead of each animal by dipping a paint brush in the mixture and making a single 15 cm stroke on the forehead. Application may also be made with a sprayer calibrated to apply 5.5 mL. Apply every morning for two weeks during peak fly season while cattle are in the barn. Thereafter apply as needed. Prepare fresh bait daily.

SHEDS, STABLES, LIVESTOCK BARNs, LOAFING SHEDS, PIGS PENS, OUTDOOR AREAS: For the control of flies, gnats, mosquitoes, roaches, fleas, ants and sowbugs. Dilute 25 mL of Sanex VAPO 20 Emulsifiable Concentrate Insecticide in 1 litre of water. Apply as a coarse wet spray at the rate of 5 L per 100 m<sup>2</sup>. Remove animals and close windows and doors. Thoroughly ventilate buildings before returning animals. Do not use in milk rooms or milk processing rooms.

POULTRY HOUSES: To control flies, gnats, mosquitoes, roaches, ants, sowbugs. Prepare spray as outlined above. Apply to window sills and outside of penned enclosures at the rate of 5 mL per 100 m<sup>2</sup>.

OUTDOOR LIVING AREAS, PICNIC GROUNDS, BACKYARD AREAS, PATIOS, LATRINES, LOADING DOCKS, PARKING AREAS, REFUSE AREA, AND OTHER AREAS AROUND BUILDINGS: To control flies, gnats, mosquitoes, roaches, fleas, ants, sowbugs. Prepare spray as outlined above. apply at the rate of 5 L per 100 m<sup>2</sup>.

AROUND DWELLINGS AND OTHER BUILDINGS: To control honeybees, paper nest wasps, hornets, and yellowjackets. Prepare a 1% spray by mixing 50 mL of Sanex Vapo 20 E.C. Insecticide in 1 litre of water. Soak areas or colonies where insects congregate. Apply at night when insects are relatively inactive.

GREENHOUSES, ORNAMENTALS, CUCUMBERS, TOMATOES: To control aphids and whiteflies. Vacate premises, close windows and doors. Spray foliage to point of run-off (approximately 5 L of spray per 100 m<sup>2</sup>) with a solution of 6 mL of Sanex Vapo 20 in 1 litre of water. Thoroughly ventilate premises before re-entering on the day following treatment. Do not apply to cucumbers or tomatoes within 7 days of harvest.

DOG KENNELS: To control flies, gnats, mosquitoes, roaches,

fleas, ants and sowbugs. Dilute 25 mL of Sanex Vapo 20 E.C. Insecticide in 1 litre of water. Treat outside dog runways, window sills, and ledges. Use 10 L per 100 m<sup>2</sup>.

PRECAUTIONS: Do not get on skin or eyes. Do not breathe fumes. Wear natural rubber gloves and protective clothing when handling. Wear a mask or respirator offering protection against dichlorvos insecticide. Keep out of reach of children. Keep in original container during storage. Do not store near nor contaminate feed or foodstuffs. Do not contaminate streams, lakes, ponds, irrigation water, water used by livestock or water used for domestic purposes. Keep product away from flame or heat.

FIRST AID: In case of skin contact, wash immediately with soap and water. Wash all contaminated clothing before re-use. In case of eye contact, flush eyes with running water for 10 minutes. Product contains petroleum distillate. If accidentally swallowed, do not induce vomiting. Keep victim still and quiet. Give artificial respiration if victim is not breathing. Call a doctor immediately.

NOTE: This product contains petroleum distillates.

TOXICOLOGY: Symptoms of poisoning include weakness, headache, tightness in chest, blurred vision, non-reactive pin-point pupils, salivations, sweating, nausea, vomiting, diarrhea, and abdominal cramps. If there are signs of parasympathetic stimulation, atropine sulphate should be injected at 10 minute intervals. In doses of 1 to 2 milligrams until complete atropinization has occurred. Morphine is contraindicated. Clear chest by postural drainage. Observe patient continuously for 48 hours. Repeated exposure to cholinesterase inhibitors may, without warning, cause prolonged susceptibility to very small doses of any cholinesterase inhibitor. Allow no further exposure until time for cholinesterase regeneration has been allowed as determined by blood test.

NON-WARRANTY: Seller's guarantee shall be limited to the terms set out on the label and subject thereto, the buyer assumes the risk to persons or property arising from the use or handling of this product and accepts the product on that condition.

NOTICE TO USER: This control product is to be used only in accordance with the directions on this label. It is an offence under the Pest Control Products Act to use a control product under unsafe conditions.

\*\*\*\*\*

This label transcript service is offered by the Pest Management Regulatory Agency to provide efficient searching for label information. This service and this information do not replace the official hard-copy label. The PMRA does not provide any

guarantee or assurance that the information obtained through this service is accurate, current or correct, and is therefore not liable for any loss resulting, directly or indirectly, from reliance upon this service.

+))