

91.03.28

POULIN'S

RODENT DOOM  
READY TO USE BAIT

COMMERCIAL

READ THE LABEL BEFORE USING

NET CONTENTS        454g

REGISTRATION NO. 14379 PEST CONTROL PRODUCTS ACT

GUARANTEE:

CHLOROPHACINONE        0.005%

CAUTION:            KEEP OUT OF REACH OF CHILDREN

POULIN EXTERMINATORS  
24 MARION PLACE,  
WINNIPEG, MANITOBA,  
R2H 0S9

To control Norway rat (*Rattus norvegicus*), and Roof rat (*Rattus rattus*) in and around farm buildings, processing plants, feed plants, feed barns, storage sheds, industrial buildings, and garbage dumps. Baits should be placed in amounts from 100 g up to 454 g in locations frequented by rats or where rats have been known to feed. Baits must be protected from pets, poultry, live-stock, etc. by bait boxes or other suitable means. Baits should be examined regularly and replenished before completely eaten. Replace bait if it moulds or decomposes.

To control mice (*Mus musculus*) follow the same general directions as for rats, except that the amount of bait placements should be smaller, 10 to 50 g every 3 to 5 meters and more placements should be made. Mice are more difficult to control than rats and complete control may take a longer period of baiting.

PROCESSING PLANTS:

(Feed, food and non-food, non-feed). In food processing plants, use only when plant is not in operation. Remove or cover all food, packaging materials and utensils before placing baits.

Remove all baits and dead rodents before re-use of the plant for food processing.

IMPORTANT:

Baiting should continue as long as evidence of feeding is observed. Except in extremely heavy infestation. This would be from five days to two weeks. Where a continuous source of infestation is present from nearby dumps or fields, permanent stations should be used and bait placed in them as often as necessary.

DISPOSAL:

Follow provincial instructions for any required additional cleaning of the container prior to its disposal. Make the empty container unsuitable for further use. Dispose of the container in accordance with provincial requirements. For information on the disposal of unused, unwanted product and the cleanup of spills, contact the regional office of Environmental Protection, Environment Canada.

PRECAUTIONS:

POULIN'S RODENT DOOM contains as its active ingredient anticoagulant chemical which if accidentally taken by humans, domestic animals or pets, this material may reduce the clotting ability of the blood and cause bleeding. In case baits are accidentally eaten, place finger at back of throat to induce vomiting and repeat until vomit fluid is clear. Call a physician immediately. Not to be used in processing areas where food may be exposed.

NOTICE TO USER:

This control product is to be used only in accordance with the directions on this label. It is an offence under the Pest Control Products Act to use a control product under unsafe conditions.

TOXICOLOGY:

Vitamin K1 in the form of intravenous, intramuscular or subcutaneous injections, or by oral ingestion are suggested remedial treatments for anticoagulant poisoning. The severity of the case measured by establishing prolonged prothrombin times (P.T.) will determine the appropriate therapy. Monitoring P.T. will indicate the necessity of repeated treatments.

\*\*\*\*\*

This label transcript service is offered by the Pest Management Regulatory Agency to provide efficient searching for label

information. This service and this information do not replace the official hard-copy label. The PMRA does not provide any guarantee or assurance that the information obtained through this service is accurate, current or correct, and is therefore not liable for any loss resulting, directly or indirectly, from reliance upon this service.

+))