

11-MAY-1998

PETCOR (TM)
FLEA SPRAY FOR DOGS + CATS
WITH PRECOR (R)
(IGR) Insect Growth Regulator
DOMESTIC

REGISTRATION NO: 22249

PEST CONTROL PRODUCTS ACT

GUARANTEE:

(S)-Methoprene.....	0.25%
Pyrethrins.....	0.18%
Piperonyl Butoxide, Technical.....	0.36%
N-Octyl Bicycloheptene Dicarboximide.....	0.60%

KEEP OUT OF REACH OF CHILDREN

READ THE LABEL BEFORE USING

CAUTION

WARNING

FLAMMABLE

EYE IRRITANT

475 mL

Wellmark International
P.O. Box 20040, Woodlawn Postal Outlet
Guelph, Ontario N1H 8H6

GENERAL INFORMATION: Kills adult cat fleas, dog fleas, American dog and brown dog ticks. Effectively breaks the flea life cycle

and prevents the growth of the pre-adult stages of the flea. One treatment kills adult fleas for up to 7 days and keep flea eggs from hatching for 30 days. Aids in control of flea bite dermatitis by killing fleas.

DIRECTIONS FOR USE:

SHAKE WELL BEFORE USING

Cover animal's eyes with hand and with a firm, fast stroke, spray head, ears and chest until damp but not soaked. With fingertips, rub into face around mouth, nose and eyes. Then spray the neck, middle and hind quarters, finishing with legs. Make sure spray thoroughly wets ticks. For best results, spray against natural lay of the hair, rub in well and, following the natural lay of the hair, wipe off excess with dry cloth and then brush. Using this technique, the application of 4 to 7 mL of Petcor Flea Spray per kilogram of body weight (i.e. 2.5 to 5 squeezes of the trigger sprayer per kilogram) will provide effective flea control.

Repeat only as necessary. Given the residual control afforded by Petcor Flea Spray, the application of this product once per month should be adequate. In exceptional circumstances, where there is a heavy external source of adult fleas re-infesting the animal, re-treatment may be increased to once per week. The use of an effective product for controlling adult and larva flea populations in the pet's environment will reduce the possibility of re-infestation of the animal.

PRECAUTIONS: KEEP OUT OF REACH OF CHILDREN. Harmful if swallowed. Avoid contact with eyes. After using, wash hands with soap and water.

Consult a veterinarian before using on pregnant animals, sick animals, or animals under drug or other pesticide treatment. Do not use on nursing animals or animals under 8 weeks of age. Discontinue use if adverse effects are observed.

Sensitive cats may exhibit temporary and non-harmful salivation when exposed to pyrethrin containing products. In these cases, the use of the lower treatment rates indicated is recommended. If condition persists, consult your veterinarian.

FIRST AID: IF SWALLOWED, call physician or Poison Control Centre. Drink 1 or 2 glasses of water and induce vomiting by touching the back of the throat with finger. Do not induce vomiting or give anything by mouth to an unconscious person. IF IN EYES, flush with plenty of water. Get medical attention if irritation persists. IF ON SKIN, wash affected area with soap

and water.

STORAGE AND DISPOSAL: Flammable. Keep away from heat or open flame. Do not reuse empty container. Wrap empty container and place in garbage.

PRECOR is a registered trademark and Pector is a trademark of Wellmark International.

E.P.A. Est. No. 7056-TX-1

This label transcript service is offered by the Pest Management Regulatory Agency to provide efficient searching for label information. This service and this information do not replace the official hard-copy label. The PMRA does not provide any guarantee or assurance that the information obtained through this service is accurate, current or correct, and is therefore not liable for any loss resulting, directly or indirectly, from reliance upon this service.

+))