

92.03.17

CLEAN CROP(R)

2,4-D
ESTER 400

COMMERCIAL

CAUTION POISON

READ THE LABEL BEFORE USING.

Controls Most Broad-Leaved Weeds In Turf & Crops

REGISTRATION NO. 3749

PEST CONTROL PRODUCTS ACT

GUARANTEE:

2,4-D 376 grams per litre isomer specific
(present as Iso Octyl ester)

UNITED AGRI PRODUCTS
R.R. #2
Dorchester, Ontario
N0L 1G5

NET: 4 L

31542

01/92

Oats: Oats are more sensitive to 2,4-D ESTER 400 and great care should be taken if the crop has to be sprayed. Use at the rate of 0.7 litres with sufficient water to cover 1 ha. Apply when oats are between 3-leaf and flag-leaf stage. Do not apply after the flag-leaf stage.

WARNING: Do not apply 2,4-D ESTER 400 if there is any danger of drift onto susceptible plants such as vegetables, flowers, etc., since even minute quantities will cause severe injury to those plants. Also, vapours of the ESTER 400 may damage susceptible plants close to the application area. Therefore, do not spray on windy or exceptionally hot days. Containers, sprayers, etc., used for 2,4-D ESTER 400 should not be used for insecticides, fungicides, or any other materials if applied to susceptible plants. Allow 24 hours between application of product and grazing/cutting of hay.

PRECAUTIONS: KEEP OUT OF REACH OF CHILDREN. Avoid contact with eyes, skin and clothing. Wash hands after use. Do not store near food, feed, fertilizer, insecticide, or fungicide. Do not contaminate streams, lakes or ponds.

FIRST AID:

If swallowed, do not induce vomiting. Call a doctor immediately. Wash affected parts thoroughly.

TOXICOLOGICAL INFORMATION: May be absorbed through the skin. Contains kerosene as a diluent.

DISPOSAL:

1. Rinse the emptied container thoroughly and add the rinsings to the spray mixture in the tank.
2. Follow provincial instructions for any required additional cleaning of the container prior to its disposal.
3. Make the empty container unsuitable for further use.
4. Dispose of the container in accordance with provincial requirements.
5. For information on the disposal of unused, unwanted product and the cleaning up of spills, contact the regional office of Conservation and Protection, Environment Canada.

NOTICE TO USER: This control product is to be used only in accordance with the directions on this label. It is an offence under the Pest Control Products Act to use a control product under unsafe conditions.

NOTICE TO BUYER: Seller's guarantee shall be limited to the terms set out on the label and subject thereto, the buyer assumes any risk to persons or property arising from the use or handling of this product and accepts the product on that condition.

MIXING DIRECTIONS: Fill your spray tank half full of water, then add 2,4-D ESTER 400 at the recommended rates, and finally add the balance of water. When mixing 2,4-D ESTER 400 with oil, add the chemical to the oil and THOROUGHLY agitate or shake for several minutes.

DIRECTIONS FOR USE:

LAWN & TURF WEED CONTROL: Use 2,4-D ESTER 400 at the rate of 30 mL per 20 litres of water to cover 100 m², or 3 litres to 3.75 litres per 2000 litres of water to 1 ha. If lawns begin to brown, water heavily. Use lower rates under rapid growing conditions, or on hot, dry days, or in dry climate. Do not use on newly seeded lawns, Bent grasses, or when there is any danger of drift. Do not spray for several days after mowing.

HARDY WEEDS: Such as Morning glory, Thistle, etc., use at the rate of 6 mL per 1 litre (600 mL per 100 litres) of water, and spray the plants thoroughly (to the point of run-off). Treat weeds when actively growing.

WOOD GROWTH OR BRUSH: Use 12 mL per 1 litre (1200 mL per 100 litres) of water, and spray to the point of run-off, make sure to obtain complete coverage. Repeat application if necessary. Spray when plants are tender and actively growing, under 2.5 m in height.

STUMP KILLING: Use 2,4-D ESTER 400 at the rate of 35 mL to 50 mL per 1 litre of water, kerosene or strained waste oil, and spray stumps and any exposed parts of roots thoroughly. Make sure to obtain complete coverage. Apply with a brush or sprayer.

CEREALS: Barley, Wheat, and Rye (not underseeded to legumes): Use 2,4-D ESTER 400 at the rate of 0.7 litres to 1.4 litres with sufficient water to give complete coverage of 1 ha. Use higher rate for more resistant weeds, heavy weed infestation, or under cooler weather conditions. Spray at the 3-leaf stage of growth. Do not spray growth underseeded to legumes unless certain damage to legumes can be tolerated.

This label transcript service is offered by the Pest Management Regulatory Agency to provide efficient searching for label information. This service and this information do not replace the official hard-copy label. The PMRA does not provide any guarantee or assurance that the information obtained through this service is accurate, current or correct, and is therefore not liable for any loss resulting, directly or indirectly, from reliance upon this service.

+))

+))