

85.12.31

DOMESTIC 300 g

WILSON

GARDEN SULPHUR

FUNGICIDE/MITICIDE

Spray or Dust

FOR FLOWERS, FRUITS, VEGETABLES
AND FOR ACIDIFICATION OF SOIL

REG. NO. 17546 P.C.P. ACT

Guaranteed Minimum Analysis:

Sulphur (S) 92%
Neutralizing Value 100%
(CaCO₃ Equivalent)

READ THE LABEL BEFORE USING

WILSON LABORATORIES INC.
DUNDAS, ONTARIO
L9H 3H3

DIRECTIONS FOR USE: Controls Powdery Mildew, Rust, Black Spot, Scab, Black Knot and certain mites.

How To Use: May be dusted directly onto plants or bulbs, or may be mixed with water for spray application.

Mixing: Add 15 mL (1 tablespoonful) to each litre of water and mix thoroughly. Agitate the mixture while spraying to avoid settling.

NOTE: 15 mL weighs approximately 10 grams.

SOIL AMENDMENT: To acidulate soil for acid loving plants, work in 15 mL (1 tablespoonful) per square metre around the plants 5 cm deep. Treat heavy clay soils in the fall and work in 15 cm

deep.

APPLES, PEARS: Powdery Mildew, Scab and Mites. Apply from green tip to first cover spray. Reduce rate to one half for subsequent sprays. Do not use on Anjou or Comice pears.

CHERRY (sweet and sour): Brown Rot, Leaf Spot, Powdery Mildew, Mites. Apply in scheduled protective spray program from bloom to harvest.

GRAPES AND STRAWBERRIES: Powdery Mildew. Apply at first sign of mildew and repeat every ten days if necessary. Do not use on Concord grapes or other sensitive varieties.

PEACHES AND PLUMS: Brown Rot, Leaf Spot and Black Knot.

CURRENTS AND GOOSEBERRIES: Powdery Mildew, Rust. Apply before earliest blossoms and continue through harvest.

BEANS, PEAS: Powdery Mildew, Rust. Spray or dust on appearance of disease and repeat at 7 to 10 day intervals as necessary.

TOMATO: Russet Mites. Spray or dust when the fruit begins to set.

ROSES, FLOWERS, ORNAMENTALS: Rust, Powdery Mildew, Black Spot, Mites. Dust or spray every 7 to 10 days through the season.

PREHARVEST INTERVAL: May be used up to one day before harvest on all crops except grapes for wine which require a 21 day interval.

PRECAUTIONS: KEEP OUT OF REACH OF CHILDREN. Do not apply at temperatures 24°C or higher. Do not apply to cucumbers, melon and squash. Do not apply within 30 days of oil sprays. Avoid breathing spray dust or mist. Avoid contact with skin, eyes, and clothing. Do not contaminate feed, food or any body of water. Wash thoroughly after use and before eating, drinking or smoking. Store in a dry ventilated place. Dispose of empty container with regular garbage.

FIRST AID: May cause irritation to skin or eyes. If on skin, wash thoroughly with soap and water. If in eyes, flush with water for 5 to 10 minutes and obtain medical attention. If swallowed, give patient 1 or 2 glasses of water and induce vomiting by inserting a finger down the throat. Obtain medical attention. Do not try to cause vomiting if the patient is unconscious.

This label transcript service is offered by the Pest Management Regulatory Agency to provide efficient searching for label information. This service and this information do not replace the official hard-copy label. The PMRA does not provide any guarantee or assurance that the information obtained through this service is accurate, current or correct, and is therefore not liable for any loss resulting, directly or indirectly, from reliance upon this service.

+))