

96.01.15

CLEAN CROP (R)

FERBAM 76WG

Fungicide

Wettable Granular

COMMERCIAL

GUARANTEE:

Ferbam..... 76%

REGISTRATION NO. 14610  
PEST CONTROL PRODUCTS ACT

READ THE LABEL BEFORE USING

NET 2 kg

United Agri Products  
R.R. #2  
Dorchester, Ontario  
N0L 1G5

DIRECTIONS FOR USE: The rates "per 1000 litres" recommended for tree fruits based on dilute spray. Ferbam 76WG may be used in low volume or as concentrate sprays by increasing the dosage to apply the same quantity of Ferbam per hectare as would normally be applied when spraying dilute mixtures. Apply in sufficient water for thorough coverage of all plant parts; this volume depends on spray equipment and size of plants.

MIXING: Ferbam 76WG mixes readily with water. Ferbam 76WG is compatible with most commonly used insecticides and fungicides except Bordeaux mixture and fixed copper plus hydrated lime.

CROP/DISEASE

APPLE: Bitter rot, black rot, brooks spot, calyx-end rot, cedar-apple rust, fly speck, frog-eye leaf spot, quince rust, scab, sooty blotch.

## METHOD/RATE

For scab apply 1.25 kg - 2.0 kg Ferbam 76WG in a regular protective schedule from green tip up to and including second cover spray. After the second cover spray reduce the rate to 1.0 kg per 1000 litres if scab is at a low level in the orchard. For the rust diseases, follow the scab program through the first cover spray and include a special bloom spray of 750 g per 1000 liters. Use the higher rates recommended for scab as long as any of the other diseases threaten. May cause russetting of Golden Delicious, Jonathan, and other sensitive varieties. Do not apply within 7 days of harvest.

PEAR: Fruit Spot, leaf blight, pear scab, sooty blotch. Apply in the pink, calyx, and first and second cover sprays using 2.0 kg FERBAM 76WG, per 1000 litres. Reduce rate to 1.25 kg in the summer sprays. Do not apply within 7 days of harvest.

APRICOT: Brown rot, coryneum blight, green rot. Apply 2.50 kg Ferbam 76WG in 1000 liters at red bud, early bloom and full bloom. Do not apply within 21 days of harvest.

PEACH: Leaf curl, coryneum blight (shot hole) Apply 1.75 kg - 3.50 kg Ferbam 76WG 1000 litres in the dormant and popcorn stages.

PEACH: brown rot, blossom blight Apply 1.75 kg - 2.25 kg Ferbam 76WG at pink bud and 25 to 75% of full bloom. Do not apply within 21 days of harvest.

PLUM, PRUNE: Brown rot Apply 2.0 - 2.50 kg Ferbam 76WG in 1000 litres at popcorn, full bloom, and 2 weeks after full bloom. Make another application 5 weeks before harvest and repeat weekly during wet weather. Do not apply within 7 days of harvest.

CERRY (SWEET, SOUR): Brown rot blossom blight, brown rot stem rot, leaf spot. For brown rot, apply 1.75 kg - 2.0 kg Ferbam 76WG in 1000 litres at pink bud, at full bloom, at petal fall, and at intervals of 7 to 10 days starting 2 to 3 weeks before harvest. For brown rot, make 3 pre-harvest applications at intervals of 14 days, beginning at petal fall, and apply again immediately after harvest. Do not apply within 4 days of harvest.

GRAPE: Black rot Apply 2.0 kg - 2.50 kg Ferbam 76WG in 1000 litres just before bloom, just after bloom, 14 days after bloom, and in supplemental applications as recommended by local authorities. Do not apply within 7 days of harvest.

BLACKBERRY, BOYSENBERRY, DEWBERRY, RASPBERRY: Anthracnose, rust, spur blight.

Apply 3.75 kg Ferbam 76WG in 1000 litres as a delayed dormant spray. Make a second application using 2.0 kg - 2.5 kg in 1000 litres when new canes are 25-30 cm tall and a third just before bloom. Do not apply to blackberry, boysenberry, dewberry or raspberry after the berries begin to form.

BLUEBERRY: Boytrytis blight, blossom blight, twig blight.

Apply 3.75 kg Ferbam 76WG in 1000 litres. Make first application when leaves are emerging or flower buds breaking. Additional applications up to mid-bloom may be made at intervals of 7-10 days, depending on the weather. Do not apply within 40 days of harvest.

TOBACCO: Blue mold (in seed beds)

Mix 5.0 kg Ferbam 76WG in 1000 litres of water and apply 20-24 litres per 100 m<sup>2</sup> when tobacco leaves are the size of a dime or when blue mold first appears. Repeat 3 times per week until transplanting.

PRECAUTION: KEEP OUT OF REACH OF CHILDREN.

Hazardous if swallowed, inhaled or absorbed through the skin. After handling, wash thoroughly with soap and water before eating, drinking or smoking and change to clean clothing. Do not apply or allow to drift to areas occupied by unprotected persons and animals or to streams, lakes or ponds to protect wildlife. Avoid contamination of feed or food, including such crops on which residue is unsafe.

FIRST AID: If swallowed, induce vomiting by sticking fingers down the throat, repeat until vomit fluid is clear. Rush patient to nearest hospital taking this label container with you. In case of skin contact, remove contaminated clothing and wash skin with soap and water. If in eyes, flush with plenty of water for 15 minutes and go to the nearest hospital at once.

STORAGE: Store product in a cool dry place--do not freeze.

DISPOSAL:

1. Follow provincial instructions for any additional cleaning of the container prior to its disposal.
2. Make the empty container unsuitable for further use.
3. Dispose of the container in accordance with provincial requirements.

4. For information on the disposal of unused, unwanted product and the cleanup of spills, contact the Provincial Regulatory Agency or the Manufacturer.

NOTICE TO USER: This control product is to be used only in accordance with the directions on this label. It is an offence under the Pest Control Products Act to use a control product under unsafe conditions.

NOTICE TO BUYER: Seller's guarantee shall be limited to the terms set out on the label and subject thereto the buyer assumes the risk to persons or property arising from the use of handling of this product and accepts the product on that condition.

\*\*\*\*\*

This label transcript service is offered by the Pest Management Regulatory Agency to provide efficient searching for label information. This service and this information do not replace the official hard-copy label. The PMRA does not provide any guarantee or assurance that the information obtained through this service is accurate, current or correct, and is therefore not liable for any loss resulting, directly or indirectly, from reliance upon this service.

+))