

15-AUG-2000

DISPAR®

TRAP-N-A-SAK II

RODENTICIDE

THE SWEET CORN BAIT RODENTS LIKE

KILLS RATS AND MICE

DOMESTIC

WITH DIPHACINONE

A PELLETTIZED RODENTICIDE "CONTAINING SWEET CORN"  
READY MIXED

USE INDOORS AND OUTDOORS

GUARANTEE: 0.005% DIPHACINONE

KEEP OUT OF REACH OF CHILDREN

READ THE LABEL BEFORE USING

6 X 85g

REG. NO. 22069 P.C.P. ACT

DISPAR DIVISION OF  
VETOQUINOL N.-A. INC.  
LAVALTRIE, (QUEBEC), CANADA J0K 1H0

INDICATIONS: For control of Norway rats (*Rattus norvegicus*), roof rats (*Rattus rattus*) and house mice (*Mus musculus*) in and around homes. Treated baits must be placed in tamper-resistant bait boxes or in locations not accessible to children, pets, domestic animals or wildlife. Do not place bait in areas where there is a possibility of contaminating food or surfaces that come in direct contact with food.

SELECTION OF TREATMENT AREAS: Determine areas where rats

and/or mice will most likely find and consume the bait. Generally, these areas are along walls, by gnawed openings, in or beside burrows, in corners and concealed places, between floors and walls, or in locations where rodents or their signs have been observed. Protect bait from rain or snow. Remove as much food as possible.

**DIRECTIONS:**

**RATS:** Apply 100 to 500 grams of bait per placement. Maintain an uninterrupted supply of fresh bait for at least 10 days.

**MICE:** Apply 7 to 15 g (1-2 tablespoons) of bait at 1-3 m intervals. Larger placements (up to 60 g) may be needed at points of very high mouse activity. Maintain an uninterrupted supply of fresh bait for at least 15 days.

**RATS AND MICE:** Replace contaminated or spoiled bait immediately. Collect and dispose of all dead animals and unconsumed bait according to local requirements. To prevent reinfestation, eliminate food, water, and harborage as much as possible. If reinfestation does occur, repeat treatment. Where a continuous source of infestation is present, establish permanent bait stations and replenish bait as needed.

**PRECAUTIONS:** KEEP OUT OF REACH OF CHILDREN. Keep away from humans, domestic animals and pets. If swallowed, this material may reduce the clotting ability of the blood and cause bleeding.

**FIRST AID:** If swallowed, give patient one to two glasses of water and induce vomiting by inserting finger down throat; repeat until vomit fluid is clear. Do not give anything by mouth to an unconscious person. Call a poison control centre immediately and contact a physician.

**TOXICOLOGICAL INFORMATION:** Vitamin K<sub>1</sub>, in the form of intravenous, intramuscular or subcutaneous injections, or by oral ingestion are suggested remedial treatments for anticoagulant poisoning. The severity of the case measured by establishing prolonged prothrombin times(P.T.) will determine appropriate therapy. Monitoring P.T. will indicate the necessity of repeated treatments.

**DISPOSAL:** Dispose of empty container in household garbage.

**NOTICE TO BUYER:**

Seller's guarantee shall be limited to the terms set out on the label and subject thereto. The buyer assumes the risk to persons or property arising from the use or handling of this product and accepts the product on that condition.

\*\*\*\*\*

This label transcript service is offered by the Pest Management Regulatory Agency to provide efficient searching for label information. This service and this information do not replace the official hard-copy label. The PMRA does not provide any guarantee or assurance that the information obtained through this service is accurate, current or correct, and is therefore not liable for any loss resulting, directly or indirectly, from reliance upon this service.

+) )