

95.09.29

MATAVEN*

AGRICULTURAL

HERBICIDE

NET CONTENTS; 10 LITRES

EMULSIFIABLE CONCENTRATE FOR CONTROL OF WILD OATS.

GUARANTEE:

FLAMPROP-METHYL.....84 GRAMS PER LITRE

READ THE LABEL BEFORE USING

REGISTRATION NO.: 14337
PEST CONTROL PRODUCTS ACT

WARNING POISON

IRRITANT

Cyanamid Crop Protection
88 McNabb St.
Markham, Ontario
L3R 6E6

NOTICE TO USER: This control product is to be used only in accordance with the directions on this label. It is an offence under the PEST CONTROL PRODUCTS ACT to use a control product under unsafe conditions.

NOTICE TO BUYER: Seller's guarantee shall be limited to the terms set out on the label and subject thereto, the buyer assumes the risk to persons or property arising from the use or handling of this product and accepts the product on that condition.

FIRST AID: This product contains petroleum distillate. If ingested do not induce vomiting. If patient is unconscious, give him air. Keep patient prone and quiet. Transport the patient immediately to the nearest physician or contact a poison control centre.

Skin Contact: If spilled on the skin, remove all contaminated clothing at once. Thoroughly wash skin with soap and water. Contact physician or contact a poison control centre. Wash

contaminated clothing before re-use.

For Eyes: If the material gets into the eyes, flush immediately with running water for at least 15 minutes. Contact physician or contact a poison control centre.

CAUTION:

KEEP OUT OF REACH OF CHILDREN. Product is harmful if swallowed. Avoid contact of this product with eyes. When handling, the use of safety goggles is recommended. Avoid inhalation of vapours. Keep product in original closed container and store away from food and feed. Wash hands and exposed skin before eating, drinking, smoking and after work. Flush concentrate from skin and eyes with water immediately after contact.

DECONTAMINATION AND DISPOSAL:

1. Rinse the emptied container thoroughly and add the rinsings to the spray mixture in the tank.
2. Follow provincial instructions for any required additional cleaning of the container prior to its disposal.
3. Make the empty container unsuitable for further use.
4. Dispose of the container in accordance with provincial requirements.
5. For information on the disposal of unused, unwanted product, and the cleanup of spills, contact the Provincial Regulatory Agency or the Manufacturer.

GENERAL INFORMATION: MATAVEN is a post-emergent herbicide for the control of wild oats in licensed varieties of wheat (spring, durum, winter), triticale, canary grass, sunflowers and certain forage crops (seed production only).

TIMING OF APPLICATION: MATAVEN can be applied to wild oats from the 3-leaf stage to the 6-leaf stage. For optimum control and maximum yield response, apply Mataven when the majority of the wild oats have 3-4 leaves.

TIME AND RATE OF APPLICATION:

CROP	CROP STAGE	LITRES MATAVEN PER HECTARE	REMARKS
------	------------	-------------------------------	---------

Wheat (licensed varieties of spring, durum, winter) Canary grass Triticale	Not beyond 6-leaf stage	5.0	
Sunflowers	Not beyond 6-leaf stage, but before dense crop canopy forms	5.0-6.5	Use the higher rate when wild oat pressure is heavy and for band application over crop row. Band application should be combined with interrow cultivation.

FORAGE CROPS FOR SEED PRODUCTION ONLY

CROP	CROP STAGE	LITRES MATAVEN PER HECTARE	REMARKS
Alfalfa, Red Clover, Birdsfoot trefoil Sanfoin, Cicer milk vetch	In year of seeding	5.0	Do not harvest or graze forages in year of treatment.
Bromegrass, Meadow Fescue, Creeping Red Fescue, Russian Wild Ryegrass, Intermediate Wheatgrass, Crested Wheatgrass	In year of seeding	5.0-7.5	Use the high rate when grown without a companion crop. Use the low rate when grown with a companion crop for which Mataven is registered. Do not harvest for forage or graze in year of treatment.

APPLICATION INSTRUCTIONS:

Ground Application: Apply Mataven in a minimum of 100 litres of water per hectare, using flat fan spray tips at a minimum pressure of 300 kPa. Use 50 mesh or larger in-line and nozzle screens. If crop or weed canopy is dense, increase water volume to improve coverage and weed control.

Aerial Application: Apply Mataven by air with not less than 15 litres per hectare of water. Do not spray in winds greater than 15 km/hr. Do not allow spray drift to contaminate other crops. Application should not be made by air when oat crops are adjacent and downwind of the treated area.

USE PRECAUTIONS:

- Allow 4 days between an MCPA or bromoxynil application and an application of MATAVEN.
- Allow 7 days between a 2,4-D or dicamba application and an application of MATAVEN.
- Rainfall within 2 hours of application may reduce effectiveness of MATAVEN.
- Protect MATAVEN from freezing.

This label transcript service is offered by the Pest Management Regulatory Agency to provide efficient searching for label information. This service and this information do not replace the official hard-copy label. The PMRA does not provide any guarantee or assurance that the information obtained through this service is accurate, current or correct, and is therefore not liable for any loss resulting, directly or indirectly, from reliance upon this service.

+))