

08-JUL-2003 Notification received of label changes. See below for these changes.

05-JUL-2004 Notification received of a label change. See below for this change.

14-APR-1998

((BOOKLET))

Notification Change

~~CLEAN CROP~~®

ESTEMINE®
2,4-D

COMMERCIAL NON-VOLATILE

LIQUID HERBICIDE

READ THE LABEL AND ATTACHED BOOKLET BEFORE USING

GUARANTEE:

2,4-D (isomer specific) 470 g/L
(present as amine salt)

REGISTRATION NO. 21667
PEST CONTROL PRODUCTS ACT

In case of spills, poisoning or fire,
telephone emergency response number
1-800-561-8273 (24 hours a day)

Notification Change

~~UNITED AGRI PRODUCTS~~
~~R.R. #2~~
~~Dorchester, Ontario~~
~~N0L 1G5~~

UNITED AGRI PRODUCTS CANADA INC. □
789 DONNYBROOK DRIVE □
DORCHESTER ON □
N0L 1G5

NET: 10 L

PRECAUTIONS: KEEP OUT OF REACH OF CHILDREN. Causes skin and eye irritation. Avoid contact with skin, eyes and clothing. Do not inhale fumes. Avoid breathing spray mist.

PROTECTIVE CLOTHING AND EQUIPMENT

Handling the concentrate (mixing and loading): wear long-sleeved shirt, long pants, chemical-resistant gloves, socks and shoes, and protective eye wear (face shield or safety glasses). Rinse gloves before removal.

Coveralls or a chemical-resistant apron should also be worn when open pouring from containers greater than 5 L.

Handling the dilute spray solution (during application or repairing or cleaning equipment): wear a long-sleeved shirt, long pants, chemical-resistant gloves, socks and shoes. Rinse gloves before removal.

Gloves are not required during application when applicator is in an enclosed tractor or an enclosed airplane cockpit.

OPERATOR USE PRECAUTIONS

- Wear freshly laundered clothing and clean protective equipment daily.
- Rinse gloves before removal.
- Wash hands before eating, drinking, using tobacco or using the toilet.
- If herbicide penetrates clothing, remove immediately, then wash thoroughly and put on clean clothing. Throw away clothing and other absorbent materials that have been drenched or heavily contaminated with his product's concentrate.
- After using the product, remove clothing and launder separately, and promptly and thoroughly wash hands and exposed skin with soap and water. Follow manufacturer's instructions for cleaning personal protective clothing and equipment. If no such instructions for washables are provided, use detergent and hot water. Keep and wash personal protective equipment separate from household laundry.
- After work, remove all clothing and shower using soap and water.

DIRECTIONS FOR USE:

CROP USE:

GROUND APPLICATION: Apply 50-200 L of spray solution per hectare depending on the type of application equipment used. Use sufficient water for even distributing. Spray at low pressures (200-275 kPa) when the weeds are actively growing.

MIXING: Shake this product well before using. Half fill the spray tank with clean water. Add the required amount of product and agitate thoroughly. Fill the tank with clean water and agitate well before use. NOTE: If spray solution has been left standing, agitate thoroughly before use.

Transfer systems for:

Containers larger than 20 Litres. Use a transfer system that avoids open pouring when transferring the liquid concentrate from such containers into the spray tank.

USE PRECAUTIONS FOR AERIAL APPLICATION (to cereal crops, grass crops, pasture rangeland, stubble and non-crop land).

Use where aerial use is appropriate suitable and safe. Exercise special care.

- . Avoid direct applications to any body of water. Do not contaminate water through spray drift or by cleaning of equipment or disposal of wastes.
- . Apply a minimum of 30L of spray solution per hectare.
- . Use boom pressure of 235 kPa or less. Avoid placing nozzles where spray will enter wing tip vortices.
- . Do not apply this product directly to, or otherwise permit it to come into direct contact with desirable crops or other desirable broadleaf plants or non-target species and do not permit spray mists to drift onto them.
- . Coarse sprays are less likely to drift, use only nozzles or nozzle configuration which minimize the production of fine spray drops. Do not angle nozzles forward into the airstream and do not increase spray volume by increasing nozzle pressure. When spraying, avoid a combination of pressure and nozzle type that will result in fine particles (mist) which are more likely to drift. A spray thickening agent or drift retardant may be used with this product to aid in reducing spray drift.
- . Do not use human flaggers.
- . Avoid Spray Drift: Apply only when there is little or no hazard from spray drift. Small quantities of the spray, which may not be visible, may seriously injure susceptible crops and damage sensitive non-target habitat. A method must be used to detect air movement, lapse conditions, or temperature inversions (stable air) such as the use of balloons or a continuous smoke column at or near the spray site or a smoke generator on the spray equipment. If the smoke develops into layers or indicates a potential for hazardous spray drift, do not spray.
- . Buffer Zones: Appropriate buffer zones should be established between treatment areas and aquatic systems and treatment areas and significant wildlife habitat.
- . Do not spray in winds exceeding 8 km per hour.
- . Do not spray in dead calm near sensitive plants. The "cloud" of suspended droplets may drift onto sensitive plants when the wind comes up. Spray only when the wind is blowing away from a sensitive crop, shelter-belt or garden.

USE PRECAUTIONS: Avoid spray drift to any desirable vegetation NOT

listed on this label as damage may occur. Avoid contamination of ponds, streams, rivers and any other water sources. Do not spray during periods of high winds when spray is likely to drift. Coarse sprays are less likely to drift. Do not spray during periods of high temperatures when crops may be under stress as damage to the crop may occur.

SPRAYER: Clean all spray equipment thoroughly before use. Select nozzle tips to apply recommended volume per hectare. Higher water volumes will reduce the risk of crop injury. Calibrate the sprayer and adjust the boom height to ensure uniform coverage. Nozzles must be in good condition and spraying the same volume. Flat fan nozzle tips are recommended.

CLEANING OF SPRAYER: Thoroughly clean the herbicide sprayer before applying and other chemical on crops which may be damaged (see PRECAUTIONS). Wash the outside of the sprayer and drain the tank completely. Remove and clean filters, screens and nozzle tips separately. Fill the sprayer tank with clean water. With the agitator running flush out the lines and boom, then drain. Fill the sprayer with clean water and use TANK and EQUIPMENT CLEANER or for each 100 L add 1 L household ammonia. Operate the pump and agitator for 15 minutes. If possible, let the solution remain in the tank and hoses overnight; recirculating and drain each time. DO NOT contaminate irrigation or drinking water when cleaning the sprayer.

TIMING: For best results spray when WEEDS ARE YOUNG AND ACTIVELY GROWING. Apply in good growing conditions. Application must be made before the crop canopy shields the weeds.

SPOT TREATMENT: For knapsack application of ESTEMINE 2,4-D for spot treatment of weeds such as thistles, mix 280 mL of product in 10 L of water. Wet all foliage thoroughly.

SELECTIVE WEED CONTROL IN SMALL GRAINS: Weeds differ in their susceptibility to ESTEMINE 2,4-D, and not all types can be controlled satisfactorily in crops. The amount of ESTEMINE 2,4-D to use will depend upon the susceptibility and whether the crops will tolerate this amount.

WHEAT, BARLEY, RYE (not underseeded with legumes): Crop may be sprayed from the 4-leaf to the early flag leaf (shot blade) stage. Do not apply after the flag leaf stage. Treatment at the 4-leaf stage usually combines maximum crop tolerance with maximum weed susceptibility. The recommended crop rate is 0.7 to 1.1 L/ha. Higher rates may be needed to control certain weed species. Possible crop injury from these higher rates is usually offset by increased yield of the cleaner crop. If applying by ground use 50 or more L/ha of water. Spraying with higher volumes of water (100-150 L/ha of water) may reduce the risk of crop injury.

PRE-HARVEST INTERVALS:

- . Do not permit lactating dairy animals to graze treated fields within 7 days after application.
- . Do not harvest forage or cut hay within 30 days after application.
- . Withdraw meat animals from treated fields at least 3 days before slaughter.

FALL RYE, WINTER WHEAT: Treat in the spring between full tillering and just before the flag leaf stage. DO NOT APPLY DURING AND AFTER THE FLAG LEAF STAGE. DO NOT apply in the fall. The recommended rate is 0.7 to 1.1 L/ha. Higher rates may be needed to control certain weed species. Possible crop injury from these higher rates is usually offset by the increased yield of the cleaner crop. If applying by ground use 50 or more L/ha of water. Spraying with higher volumes of water (100 - 150 L/ha) may reduce the risk of crop injury.

PRE-HARVEST INTERVALS:

- . Do not permit lactating dairy animals to graze treated fields within 7 days after application.
- . Do not harvest forage or cut hay within 30 days after application.
- . Withdraw meat animals from treated fields at least 3 days before slaughter.

TANK MIX: Consult the product labels to be tank mixed with ESTEMINE 2,4-D for detailed information regarding weeds controlled, precautions, warnings, restrictions, tank mixing and spraying instructions.

ESTEMINE 2,4-D PLUS TCA

For control of both green foxtail and 2,4-D susceptible broadleaf weeds in barley, a tank mix of ESTEMINE 2,4-D plus TCA may be used.

Crop	Rate of Application (Product/ha) ESTEMINE 2,4-D + TCA	Minimum Water Volume (litres/ha)	Crop Leaf Stage
Barley	700 mL - 1.1 L + 1.3 kg	100	4 - leaf

Note:

- (1) Do not apply TCA to barley seeded less than 5-7 cm deep.
- (2) Application of TCA should be made at the 1 to 3-leaf stage of green foxtail.

Notification Change

~~ESTEMINE 2,4-D PLUS BAR-FOX~~

~~For control of green foxtail and ESTEMINE 2,4-D susceptible broadleaf weeds in barley, a tank mix of ESTEMINE 2,4-D plus BAR-FOX may be used.~~

Crop	Rate of Application (Product/ha) ESTEMINE 2,4-D + BAR-FOX	Minimum Water Volume (litres/ha)	Crop Leaf Stage
Barley	700 mL - 1.1 L + 1.3 kg	100	4 leaf

ESTEMINE 2,4-D PLUS BANVEL

Crop	Rate of Application (Product/ha) ESTEMINE 2,4-D + BANVEL	Minimum Water Volume (litres/ ha)	Crop Leaf Stage
Spring Wheat	700 - 850 mL + 230 - 290 mL	100	2 - 5 leaf
Winter Wheat	700 - 850 mL + 230 - 290 mL	100	Apply in the spring when crop is 15-25 cm tall (or before shot blade stage)
Barley	700 - 850 mL + 230 mL	100	2 - 3 leaf

ESTEMINE 2,4-D PLUS PARDNER

For control of wild buckwheat, tartary buckwheat, annual smartweeds, cow cockle, volunteer sunflowers, common groundsel and ESTEMINE 2,4-D susceptible broadleaf weeds in spring wheat and barley, a tank mix of ESTEMINE 2,4-D plus PARDNER may be used.

Crop	Rate of Application (Product/ha) ESTEMINE 2,4-D + PARDNER	Minimum Water Volume (litres/ha)	Crop Leaf Stage
Spring Wheat Barley	550 - 850 mL - 1.25 L	100	4-leaf to early flag leaf

ESTEMINE 2,4-D PLUS SENCOR 500F

For control of hempnettle, chickweed, corn spurry, annual smartweeds, volunteer rapeseed, night-flowering catchfly, tartary buckwheat, common groundsel, henbit and ESTEMINE 2,4-D susceptible broadleaf weeds in spring wheat and barley, a tank mix of ESTEMINE 2,4-D plus SENCOR 500F may be used.

Crop	Rate of Application (Product/ha) ESTEMINE 2,4-D + SENCOR 500F	Minimum Water Volume (litres/ ha)	Crop Leaf Stage
Spring Wheat	850 mL - 1.1 L + 275 mL - 425 mL	100	3 - 5 leaf (mid tillering)
Barley	850 mL - 1.1 L + 275 mL - 550 mL	100	3 - 5 leaf (mid tillering)

Note: Do not apply ESTEMINE 2,4-D plus SENCOR tank mix combinations to Klondike barley.

Notification Change

~~ESTEMINE MCPA PLUS TORCH D.S.~~

~~For control of wild buckwheat, Tartary buckwheat, annual smartweeds, cow cockle, knawel, night-flowering catchfly, common groundsel and ESTEMINE 2,4-D susceptible broadleaf weeds in spring wheat, and barley, a tank mix of ESTEMINE 2,4-D plus TORCH D.S. may be used.~~

Crop	Rate of Application (Product/ha) ESTEMINE 2,4-D + TORCH D.S.	Minimum Water Volume (litres/ha)	Crop Leaf Stage
Spring Wheat Barley	550 - 850 mL - 625 L	100	4 - leaf to early flag leaf

FIELD CORN: Apply as an overall spray before the corn is 15 cm tall (leaf extended) and/or before the 6-leaf stage. DAMAGE TO THE CORN MAY OCCUR IF AN OVERALL SPRAY IS APPLIED BEYOND THIS STAGE.

The recommended crop rate for an overall spray is 0.6 m to 1.2 L/ha. Do not exceed this crop rate for an overall spray. For later applications (corn 15-75 cm leaf extended) a shielded spray or directed spray with drop pipes is necessary. Keep spray off the corn foliage. DO NOT apply within 2 weeks of silking and tasselling. Use 100 or more L/ha of water. In case of very dry conditions, delay spray until one

week following rain. Avoid cultivation for 2 weeks following spraying. DO NOT add oil or surfactant.

PRE-HARVEST INTERVALS:

- . Do not permit lactating dairy animals to graze fields within 7 days after application.
- . Do not harvest as forage within 30 days after application.
- . Withdraw meat animals from treated fields at least 3 days before slaughter.

GRASS CROPS - FOR FORAGE AND SEED PRODUCTION

Seedling Stage - Stand Establishment. May be used on seedling grasses from the 3-leaf to just before the shot-blade of development. Spray thoroughly with .585 to 1.1 L/ha of product in 100 to 200 L water.

Established Grasses: May be used in the spring up to the shot-blade of the grass or in the fall after harvest to control annual and biennial broad-leaved weeds. On seed crops do not exceed 1.1 L/ha of product. On established stands of forage grasses grown for hay or pasture, spray thoroughly using 0.7 to 2.3 L/ha of product in 100 to 300 L water.

Late germinating weeds such as red bartsias may be controlled at 1.1 L/ha of product after the first hay crop has been removed.

Do not use during flower development or pollination or seed yields will be reduced.

PRE-HARVEST INTERVALS:

- . Do not permit lactating dairy animals to graze fields within 7 days after application.
- . Do not harvest forage or cut hay within 30 days after application.
- . Withdraw meat animals from treated fields at least 3 days before slaughter.

ESTABLISHED PASTURES AND RANGELAND (without legumes): Apply at the time of rapid growth (usually May, June and/or September). Spray thoroughly using 1.5 to 2.75 L/ha of ESTEMINE 2,4-D with 100 to 300 L of water. (Use the higher rate for perennial weeds, a second treatment may be required for spot spraying). Apply only after pasture has been grazed or cut and weed growth has resumed.

There is a possibility that plants poisonous to livestock will appear more palatable and could be grazed more readily after a 2,4-D treatment (eg: tansy ragwort, water hemlock poison hemlock, death camas).

Suitable precautions should be taken when such plants are present. Livestock should not be allowed to graze the treated area until poisonous plants have been removed, are dead or have resumed a normal growth habit. Under conditions of good growth and adequate soil moisture, sprays can be applied up to 2 weeks before normal frost time.

PRE-HARVEST INTERVALS:

- . Do not permit lactating dairy animals to graze treated fields within 7 days after application.
- . Do not harvest forage or cut hay within 30 days after application.
- . Withdraw meat animals from treated fields at least 3 days before slaughter.

ASPARAGUS (Eastern Canada Only): Apply early in the season immediately after a thorough cutting or disking. The recommended crop rate is 2.75 L/ha. If emerging are contacted directly by spray drift they may become distorted ie: twisted. Spears emerging afterwards will be normal. ESTEMINE 2,4-D may also be applied 3 weeks after a post harvest disking with a directed spray to avoid the fern. Apply 2 to 2.75 L/ha in 50 to 200 L of water by ground application.

STRAWBERRIES (Eastern Canada only): Apply 2 to 4 weeks AFTER PLANTING, but before the runners root in June. Use 1.1 L/ha in 100 to 200 L of water by ground application. A low pressure nozzle and coarse spray are recommended to reduce drift.

Renovation Treatment: Apply 2 L/ha in 100 to 200 L of water after harvest but before mowing. Delay mowing for a few days after application. DO NOT apply between mid-August and fall dormancy because of possible damage to flower bud formation. Veestar is more sensitive than Redcoat. This treatment may cause temporary distortion of leaves.

RASPBERRIES (Eastern Canada only): Apply at any time EXCEPT DURING BLOOM. Avoid spraying new shoots. Apply 1.1 L/ha in 100 to 200 L of water by ground application. Higher rates may be used for spot treatments.

NON CROP USES:

GROUND APPLICATION: Apply 50-300 L of spray solution per hectare depending on the type of application system being used. Use sufficient water to ensure uniform coverage.

AERIAL APPLICATION (STUBBLE AND UNCROPPED LAND ONLY). (including fixed and rotary wing aircraft) Apply a minimum of 30 L of spray solution per hectare. Use boom pressures of 235 kPa or less. Avoid placing nozzles where spray will enter wing tip vortices. (see use precautions for aerial application)

STUBBLE LAND, ROADSIDES, UNCROPPED LAND: DO NOT APPLY BY AIR TO ROADSIDES. Apply at the time of rapid growth (usually May, June and/or September). Spray thoroughly using 1.7 to 4.5 L/ha of product in 100 L or more of water if applying by ground (Use the higher rate for perennial weeds and susceptible woody growth, a second treatment may be required as a spot treatment). Cultivating grain stubble and spraying after regrowth may aid in control. DO NOT apply on freshly seeded grasses and pastures until established.

There is a possibility that plants poisonous to livestock will appear more palatable and could be grazed more readily after 2,4-D treatment (eg: tansy ragwort, water hemlock, poison hemlock, death camas). Suitable precautions should be taken when such plants are present. Livestock should not be allowed to graze the treated area until poisonous plants have been removed, are dead or have resumed a normal growth habit. Under conditions of good growth and adequate soil moisture, sprays can be applied up to 2 weeks before normal frost time.

PRE-HARVEST INTERVALS:

- . Do not permit lactating dairy animals to graze treated fields within 7 days after application.
- . Do not harvest forage or cut hay within 30 days after application.
- . Withdraw meat animals from treated fields at least 3 days before slaughter.

LAWNS, PARKS, GOLF COURSES: Apply by ground only. Apply at time of rapid growth (usually May, June and/or September) before grasses are in the flag leaf (shot blade) stage. Apply 1.5 to 2.75 L/ha of ESTEMINE 2,4-D in 100 to 200 L of water/ha by ground application. DO NOT USE ON BENTGRASS OR NEWLY SEED LAWNS. Avoid mowing for several days before and after treatment. Do not spray within 24 hours of rainfall or during hot, dry periods.

Use caution when spraying near sensitive, desirable vegetation as contact with spray drift will cause damage.

MAXIMUM APPLICATION FREQUENCY TO TURF: For good turf/lawn management normally two applications per year per treated site are adequate. This does not include spot treatments.

RE-ENTRY INTERVAL:

Do not allow people (other than applicator) or pets on treatment area during application. Do not enter treated areas until spray has dried.

WEEDS CONTROLLED

SUSCEPTIBLE easy-to-control weeds in the seedling stage (2 to 4 leaf) and growing rapidly. Use 0.7 to 1.1 L/ha. Use the higher rate for larger weeds, dry or cold weather or heavy infestations.

Bluebur, burdock (before the 4-leaf stage), cocklebur, common plaintain, daisy fleabane, false flax, false ragweed, flixweed, goat's beard, kochia, lamb's-quarters (including triazine types), mustards (except dog and tansy), prickly lettuce, ragweeds (common and giant - including triazine resistant types), redroot pigweed (including triazine resistant types), Russian pigweed, Russian thistle, shepherd's purse, stinging nettle, stinkweed, sweet clover, thyme-leaved spurge, wild radish, wild (prairie) sunflower.

LESS SUSCEPTIBLE harder-to-control weeds in the seedling stage (2 to 4 leaf) and growing rapidly. Use 1.2 to 1.8 L/ha. Use the higher rate for larger weeds, dry or cold weather, or heavy infestations.

Weeds listed under susceptible plus annual sowthistle, blue lettuce, biennial wormwood, common chickweed, common purslane, curled dock (before the 4-leaf stage), dandelion in lawns, dog and tansy mustard, oak-leaved goosefoot, groundsel, gumweed, hairy galinsoga, hawkweed, heal-all, hedge bindweed, knotweed (before the 4-leaf stage) leafy spurge, lady's thumb, peppergrass, pineappleweed, prostate pigweed, purslane, Russian knapweed, sheep sorel, smartweed, tumble pigweed, velvet leaf, volunteer canola and mustard, wild buckwheat, yellow rocket (before the 4-leaf stage).

TOP GROWTH CONTROL ONLY: Use 1.2 to 1.8 L/ha. Biennial wormwood, blue lettuce, bull thistle, burdock, buttercup, Canada thistle, field bindweed, field dandelion, gum weed, hedge bindweed, hoary cress, leafy spurge, mouse-eared chickweed, perennial sowthistle, Russian knapweed, tartary buckwheat, teasel, volunteer sunflower, wild buckwheat, yellow rocket. Only rates greater than 1.8 L/ha will suppress top growth of hemp nettle and horsetail.

FIRST AID:

If swallowed, INDUCE VOMITING. Rush patient to nearest hospital taking this label with you or contact a poison control centre. Never give anything by mouth to an unconscious person.

In case of skin contact, remove contaminated clothing and wash skin thoroughly with soap and water.

In case of eye contact, flush eyes with plenty of water for at least 15 minutes and get medical attention or contact a poison control centre.

If inhaled, remove to fresh air and get medical attention.

STORAGE: Store container tightly closed and away from seeds, feeds, fertilizer, plants and foodstuffs. Protect from freezing. Product

exposed to subfreezing temperatures should be warmed to 5'C and mixed thoroughly before using. Insecticides and fungicides should be segregated from herbicides so as to prevent the possibility of cross-contamination. SHAKE WELL BEFORE USING.

RETURNABLE-REFILLABLE CONTAINERS

After use, return the container to the point of purchase or designated locations. This container must only be refilled with ESTEMINE 2,4-D. Do not reuse container for any other purpose. Prior to refilling, inspect thoroughly for damage such as cracks, punctures, abrasions, and damaged or worn out threads on closure devices. Check for leaks after refilling and before transportation. Do not refill or transport damaged or leaking containers. If the container is not being refilled, return to point of purchase. For information on the disposal of unused, unwanted product and the clean-up of spills contact the provincial regulatory agency or the manufacturer.

DISPOSAL:

1. Rinse the emptied container and add the rinsings to the spray mixture in the tank.
2. Follow provincial instructions for any required additional cleaning of the container prior to its disposal.
3. Make the empty container unsuitable for further use.
4. Dispose of the container in accordance with provincial requirements.
5. For information on the disposal of unused, unwanted product and the cleanup of spills contact the Provincial Regulatory Agency or the manufacturer.

NOTICE TO USER: This control product is to be used only in accordance with the directions on this label. It is an offence under the Pest Control Products Act to use a control product under unsafe conditions.

NOTICE TO BUYER: Seller's guarantee shall be limited to the terms set out on the label and subject thereto, the buyer assumes the risk to persons or property arising from the use or handling of this product and accepts the product on that condition.

®ESTEMINE AND BAR-FOX are registered trademarks of NOVARTIS CROP PROTECTION Inc.

BANVEL is a registered trademark of BASF Canada Inc.

PARDNER AND TORCH D.S. are registered trademarks of RHONE-POULENC

SENCOR is a registered trademark of BAYER AG GERMANY

((LABEL))
14-APR-1998

Notification Change

~~CLEAN CROP (R)~~

ESTEMINE®
2,4-D

COMMERCIAL

HERBICIDE

READ THE LABEL AND ATTACHED BOOKLET BEFORE USING

GUARANTEE:

2,4-D (isomer specific) 470 g/L
(present as amine salt)

REGISTRATION NO. 21667
PEST CONTROL PRODUCTS ACT

Notification Change

~~UNITED AGRI PRODUCTS~~
~~R.R. #2~~
~~Dorchester, Ontario~~
~~N0L 1G5~~

UNITED AGRI PRODUCTS CANADA INC. □
789 DONNYBROOK DRIVE □
DORCHESTER ON □
N0L 1G5

NET: 10 L

PRECAUTIONS: KEEP OUT OF REACH OF CHILDREN. Causes skin and eye irritation. Avoid contact with skin, eyes and clothing. Do not inhale fumes. Avoid breathing spray mist.

PROTECTIVE CLOTHING AND EQUIPMENT

Handling the concentrate (mixing and loading): wear long-sleeved shirt, long pants, chemical-resistant gloves, socks and shoes, and protective eye wear (face shield or safety glasses). Rinse gloves before removal.

Coveralls or a chemical-resistant apron should also be worn when open pouring from containers greater than 5 L.

Handling the dilute spray solution (during application or repairing or cleaning equipment): wear a long-sleeved shirt, long pants, chemical-resistant gloves, socks and shoes. Rinse gloves before removal.

Gloves are not required during application when applicator is in an enclosed tractor or an enclosed airplane cockpit.

OPERATOR USE PRECAUTIONS

- Wear freshly laundered clothing and clean protective equipment daily.
- Rinse gloves before removal.
- Wash hands before eating, drinking, using tobacco or using the toilet.
- If herbicide penetrates clothing, remove immediately, then wash thoroughly and put on clean clothing. Throw away clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate.
- After using the product, remove clothing and launder separately, and promptly and thoroughly wash hands and exposed skin with soap and water. Follow manufacturer's instructions for cleaning personal protective clothing and equipment. If no such instructions for washables are provided, use detergent and hot water. Keep and wash personal protective equipment separate from household laundry.
- After work, remove all clothing and shower using soap and water.

FIRST AID:

If swallowed, INDUCE VOMITING. Rush patient to nearest hospital taking this label with you or contact a poison control centre. Never give anything by mouth to an unconscious person.

In case of skin contact, remove contaminated clothing and wash skin thoroughly with soap and water.

In case of eye contact, flush eyes with plenty of water for 15 minutes and get medical attention or contact a poison control centre.

If inhaled, remove to fresh air and get medical attention or contact a poison control centre.

STORAGE: Store container tightly closed and away from seeds, feeds, fertilizer, plants and foodstuffs. Protect from freezing. Product exposed to subfreezing temperatures should be warmed to 5°C and mixed thoroughly before using. Insecticides and fungicides should be segregated from herbicides so as to prevent the possibility of cross-contamination. SHAKE WELL BEFORE USING.

DISPOSAL:

1. Rinse the emptied container thoroughly and add the rinsings to the spray mixture in the tank.
2. Follow provincial instructions for any required additional cleaning of the container prior to its disposal.

3. Make the empty container unsuitable for further use.
4. Dispose of the container in accordance with provincial requirements.
5. For information on the disposal of unused, unwanted product and the cleanup of spills, contact the Provincial Regulatory Agency or the Manufacturer.

NOTICE TO USER:

This control product is to be used only in accordance with the directions on this label. It is an offence under the Pest Control Products Act to use a control product under unsafe conditions.

NOTICE TO BUYER:

Seller's guarantee shall be limited to the terms set out on the label and subject thereto. The buyer assumes the risk to persons or property arising from the use or handling of this product and accepts the product on that condition.

This label transcript service is offered by the Pest Management Regulatory Agency to provide efficient searching for label information. This service and this information do not replace the official hard-copy label. The PMRA does not provide any guarantee or assurance that the information obtained through this service is accurate, current or correct, and is therefore not liable for any loss resulting, directly or indirectly, from reliance upon this service.

+))