

97.05.30

List No. 1023

(ALL WEATHER)

BAR BAIT (R)

Rat and Mouse Killer

(Contains Prolin)

Commercial

454 g

REGISTRATION NO. 14203

PEST CONTROL PRODUCTS ACT

GUARANTEE:

WARFARIN	0.025%
SULFAQUINOXALINE	0.025%

Solvay Animal Health, Inc.
Kitchener, Ontario
N2C 1L4

AREAS OF USE: For use in dwellings, farm buildings, food service establishments, empty granary bins, processing plants (feed, food, non-food) and storage areas (feed, food, non-food).

READ THE LABEL BEFORE USING: Read complete directions, cautions and warning statements on the back of this label.

PRECAUTION: Keep away from humans, domestic animals and pets. If swallowed, this material may reduce the clotting ability of the blood and cause bleeding.

KEEP OUT OF REACH OF CHILDREN

EPA Est. No. 61282-WI-1

DIRECTIONS FOR USE

For areas not directly related to food processing: Use only in non-food or non-feed areas where feed, food, packaging and

handling equipment are never opened or exposed.

For areas where feed or food is processed, served or stored: In meat and food processing plants (processing areas) use only when plant is not in operation. Remove or cover all food, packaging material and utensils before placing bait in baiting stations. Remove all baits and dead rodents before re-use of the plant (processing areas include storage and service).

FOR RATS: Cut through cellophane and label and break off the desired amount to be used. Assure an uninterrupted supply of bait for a period of not less than 10 days and continue baiting until all signs of feeding have stopped.

In moderately infested locations, at least 2 or more sections of this bar should be placed where rats feed, water, or travel - along walls, in corners, and concealed locations where they can feed unobserved. Each location should contain enough bait so that some will be left over the following morning.

Daily inspection is essential, especially in the first week, and bait supplies must be replenished as often as necessary. Under normal conditions, bait consumption will be light the first night, much heavier the 2nd and 3rd night thereafter. A gradual tapering-off will occur as more and more rodents perish. When no bait is being eaten (between 10 and 14 days) and signs of rodent activity have disappeared, it may be assumed that the rats in the particular location have been destroyed.

FOR MICE: Break a bar section into 4 pieces and place where mice or signs of mice have been seen. Bait stations at intervals of 2.5 to 3 m are recommended. Assure an uninterrupted supply of bait for a period of not less than 15 days.

PRECAUTIONS: Protect children, pets, wildlife and domestic animals from the bait. Harmful if swallowed. Do not expose these bait bars in areas accessible to dogs, children, wildlife and domestic animals except when enclosed within a tamper-proof bait box.

FIRST AID: If swallowed, give patient one or two glasses of water and induce vomiting by inserting finger down throat. Repeat until vomit fluid is clear. Call a physician or Poison Control Centre.

TOXICOLOGY: Vitamin K1 in the form of intravenous, intramuscular or subcutaneous injections, or by oral ingestion are suggested remedial treatments for anticoagulant poisoning. The severity of the case measured by establishing prolonged prothrombin times (P.T.) will determine appropriate therapy. Monitoring P.T. will indicate the necessity of repeated treatments.

STORAGE AND DISPOSAL: Do not contaminate water, food or feed by storage or disposal.

STORAGE: Store only in original container, in a dry place inaccessible to children and pets.

DISPOSAL:

1. Dispose of stale bait and dead rodents in accordance with local requirements.
2. Follow provincial instructions for any required additional cleaning of the container prior to its disposal.
3. Make the empty container unsuitable for further use.
4. Dispose of the container in accordance with provincial requirements.
5. For information on the disposal of unused, unwanted product and cleanup of spills, contact the Provincial Regulatory Agency or the Manufacturer.

NOTICE TO USER: This control product is to be used only in accordance with the directions on this label. It is an offence under the Pest Control Products Act to use a control product under unsafe conditions.

This label transcript service is offered by the Pest Management Regulatory Agency to provide efficient searching for label information. This service and this information do not replace the official hard-copy label. The PMRA does not provide any guarantee or assurance that the information obtained through this service is accurate, current or correct, and is therefore not liable for any loss resulting, directly or indirectly, from reliance upon this service.

+))