

95/2000

CLEAN CROP®

AZINPHOS-M® 50W

INSECTICIDE

FOR INSECT CONTROL ON FIELD, FRUIT, VEGETABLE AND ORNAMENTAL CROPS

GUARANTEE:

Azinphos-methyl ..... 50%

Contient de l'azinphos-méthyl Pour réprimer les insectes dans champs,  
fruits,

légumes et plantes ornementales

GARANTIE:

azinphos-méthyl ..... 50%

Store in cool dry place. Protect from heat.

Keep away from open flame. Do not heat.

REGISTRATION NO. 22087 PEST CONTROL PRODUCTS ACT

A emmagasiner dans un endroit frais et sec. Mettre à l'abri de la  
chaleur.

L'éloigner des flammes nues.

Ne pas le chauffer.

No. d'enregistrement 22087 Loi sur les Produits Antiparasitaires

DANGER POISON

KEEP OUT OF REACH OF CHILDREN

CALL A DOCTOR IN CASE OF ACCIDENT

GARDER HORS DE LA PORTEE DES ENFANTS

APPELER UN MEDECIN EN CAS D'ACCIDENT

READ THE LABEL BEFORE USING

LIRE ATTENTIVEMENT L'ETIQUETTE AVANT L'EMPLOI

UNITED AGRI PRODUCTS  
R.R. #2  
Dorchester, Ontario  
N0L 1G5

NET: 2 kg

PRECAUTIONS: Do not use on other crops used for food or forage. Treated orchards and groves should not be pastured within 21 days of treatment. Use only according to label directions. Application at rates above those shown may result in illegal crop residues. Do not use on food crops grown in the greenhouse. To protect fish and wildlife, do not contaminate streams, lakes, or ponds.

This product is highly toxic to bees exposed to direct treatment or residues on crops. Do not apply when crop is in bloom or allow spray to drift towards beehives. Protective information may be available from your local government extension specialist.

PRECAUTIONS: Ne pas utiliser ce produit pour traiter d'autres cultures alimentaires ou fourragères. Attendre au moins 21 jours après l'application avant de laisser paître le bétail dans les vergers et les plantations traitées. Toujours suivre les directives au pied de la lettre. Une pulvérisation plus fournie que celle qui est recommandée pourrait se traduire par des résidus illégaux dans les récoltes. Ne pas utiliser ce produit pour traiter les récoltes en serres. Afin de protéger les poissons et la faune, user des précautions nécessaires pour ne pas contaminer les rivières, les lacs et les étangs.

Ce produit se révèle très toxique pour les abeilles qui sont exposées à la pulvérisation ou aux résidus sur les récoltes. Ne faites pas de traitement quand les récoltes. Ne faites pas de traitement quand les récoltes sont en fleurs et veillez à ce que la pulvérisation ne dérive pas vers les ruches. Consultez votre biologiste-entomologiste avertisseur pour les mesures protectrices à adopter.

NOTICE TO USER: This control product is to be used only in accordance with the directions on this label. It is an offence under the PEST CONTROL PRODUCTS ACT to use a control product under unsafe conditions.

#### RESTRICTED USES

NATURE OF RESTRICTION: This product is to be stored or displayed separate from feed or food.

#### DIRECTIONS FOR USE

MIXING: AZINPHOS-M 50W mixes easily with water to form a suspension which is readily usable in all conventional ground spray equipment. Mix the required amount of AZINPHOS-M 50W with a small quantity of water. Add this premix through the screen while filling the sprayer tank or fill the tank to the required level and add the premix. Operate the agitator while mixing.

DOSAGE: Use specified dosage of AZINPHOS-M 50W in the amount of water necessary to give complete coverage of foliage. The type of equipment used will determine the concentration required.

SPRAYING: Work to windward. Protect sprayer operators from drift or mist. When low volumes of spray are applied, complete coverage and thorough application are essential for most effective results. Schedule applications in accordance with local conditions. Consult your local agricultural authorities for specific use information.

Equipment for use in spraying RAPE, SUGAR BEETS and SUNFLOWERS: AZINPHOS-M 50W may be applied using any commercial tractor drawn or self-propelled field sprayer provided it is equipped with the following:

1. Nozzle tips no finer than 6502, 8002 or TK2 with nozzle screens no finer than 0.3 mm. The above tips will provide 100L/ha when operated at 8 km/hr and 200kPa.
2. 0.3 mm or larger line strainers or screens.

NOTE: Felt filters, smaller nozzle tips or smaller screens will become clogged when using AZINPHOS-M 50W.

#### RECOMMENDED APPLICATIONS

CROP	INSECT	DOSAGE AZINPHOS-M 50W	REMARKS
FIELD CROPS Alfalfa Clover	Alfalfa plant bug, Alfalfa weevil, Aphids, Clover leaf weevil, Fleahopper, Leafhoppers, Lesser clover leaf weevil, Lygus bugs, Mites, Spittlebug	1.1 to 1.75 kg/ha	Apply specified dosage per hectare using a minimum of 80 L of water per hectare when applying with ground equipment and a minimum of 40 L per hectare in air applications. It may be necessary to use 150-200L of water per hectare on heavy growth in controlling the alfalfa weevil. Do not apply more than once per cutting, not within 21 days of harvest.

	Sweet clover weevil	1.1 kg/ha	For sweet clover weevil control, 2 applications at 7-14 day intervals may be required. Do not apply more than twice per season nor within 21 days of harvest.
Barley, Oats, Rye, Wheat	Cereal leaf beetle	850 g to 1.1 kg/ha	Apply specified dosage per hectare in sufficient water for complete coverage. Do not apply more than once per season. Do not harvest for food, feed, or forage within 30 days of treatment.
Rape	Diamondback moth	275 to 550 g/ha	Apply specified dosage per hectare using a minimum of 80 L of water. Repeat as necessary. Do not apply within 30 days of harvest. See DIRECTIONS FOR USE.
	Flea Beetles	150 to 275 g/ha	
	Red Turnip Beetle	See Remarks	Apply as a border or barrier treatment in a 18 to 30.5m wide band around the field at the rate of 27.5 to 42.5 g AZINPHOS-M 50W/1000m <sup>2</sup> . If an overall field application is needed, apply AZINPHOS-M 50W at 275 to 425 g/ha. Apply specified dosage per hectare using a minimum of 80 L of water. Repeat as necessary. Do not apply within 30 days of harvest. See DIRECTIONS FOR USE.
Sugar Beets	Flea Beetles	150 g/ha	Apply specified dosage per hectare using a minimum of 80 L of water. Do not apply more than once per season nor within 100 days of harvest. See DIRECTIONS FOR USE.

Sunflowers	Sunflower Beetles (adult)	550 g/ha	Apply specified dosage per hectare using a minimum of 80 L of water. Application can be made from the seedling stage until the sunflowers have reached full height as may be required. Two applications at 7 to 14 day intervals may be necessary. Do not apply more than 2 times per season or after heads begin to form. See DIRECTIONS FOR USE.
Tobacco	Aphids, Tobacco hornworm, Tobacco flea beetle	1.75 kg/ha	Apply specified dosage per hectare. Treat for hornworm as soon as egg masses or worms are first found. Second brood hornworms and flea beetles may require 2 applications. Repeat as necessary but not within 6 days of harvest. Prime before treating.
SEED FIELD CROPS Vetch	Aphids, Bruchid	1.1 kg/ha	Apply specified dosage per hectare. Repeat as necessary. Do not feed treated forage.

<p>FRUIT Apples, Crab Apples, Pears, Quinces</p>	<p>Aphids, Apple maggot, Codling moth, Eye-spotted budmoth, Forbes scale, Fruit tree leaf roller, Green fruitworm, Leafhoppers, Mealybug, Mites, Mullein bug (campyloma), Oblique- banded leaf roller, Oystershell scale, Pale apple leaf roller, Pear Psylla, Plum curculio, Putnam scale, Red-banded leaf roller, San Jose scale, Stink bug, Tarnished plant bug, Winter Moth</p>	<p>625 to 775 g/1000 L</p>	<p>Apply specified dosage in 1000 L of water as a full coverage spray.* For Cherries: Repeat as necessary. Do not apply more than 8 times per season nor within 15 days of harvest. For Apples, Crab Apples, Pears, and Quinces: Make 6 applications at rates of 625 to 775 g/1000 L followed by 2 applications at the 625 g/1000 L rate but not within 7 days of harvest. Consult your local agricultural authorities for proper timing for oblique-banded leaf roller on pears. AZINPHOS-M 50W is compatible with dormant and summer oils which may be added to apple and pear sprays in accordance with local recommendations. Do not use oils within 30 days of sulfur or lime-sulfur treatments. In Nova Scotia: AZINPHOS-M 50W may be used at rates of 60 to 125 g/1000 L to control pale apple leaf roller and winter moth. Consult your local agricultural authorities for specific directions on timing of applications.</p>
<p>Cherries</p>	<p>Cherry fruit fly, Forbes scale, Eye- spotted bud moth, Fruit tree leaf roller, Mites, Plum curculio, San Jose scale</p>	<p>625 g to 1.25 kg/1000 L</p>	

	Lesser peach tree borer	12.5 kg/1000 L
--	-------------------------	----------------

Apricots Peaches	Aphids-Cottony peach scale, European fruit lecanium scale, Forbes scale, Mites, Oriental fruit moth, Oblique-banded leaf roller, Platynota flavedana leaf roller, Plum curculio, Red-banded leaf roller, San Jose scale, Stink bug, Tarnished plant bug, Terrapin scale, Walnut scale, White peach scale,	625 to 775 g/1000 L	Apply specified dosage in 1000 L of water as a full coverage spray.* Repeat as necessary. Do not apply more than 8 times per season, nor within 21 days of harvest for apricots and peaches, nor within 15 days of harvest for plums and prunes. For scale control, apply when crawlers are present. Consult your local agricultural authorities for proper timing for oblique-banded leaf roller on peaches. AZINPHOS-M 50W is compatible with dormant and summer oils which may be added to peach and prune sprays in accordance with local recommendations. Do not use oils within 30 days of sulfur or lime-sulfur treatments.
	Peach twig borer	625 to 925 g /1000 L	

Plums, Prunes	Aphids, Eye-Spotted bud moth, Forbes scale, Fruit tree leaf roller, Mites, Orange tortrix, Peach twig borer, Plum curculio, Red-banded leaf roller, San Jose scale, Stink bug, Tarnished plant bug, Tussock moth,	625 g to 1.25 kg /1000 L	
	American plum borer, Peach tree borer, Lesser peach tree borer	1.25 kg/ 1000 L	
Blueberr ies	Fruitworms, Lecanium scale, Plum curculio	1.1 to 2.25 kg/ha	Apply specified dosage per hectare using approximately 1600 L of water for good coverage. Repeat as necessary. Do not apply within 14 days of harvest.
	Blueberry maggot	2.25 kg/ha	
Cranberr ies	Cranberry fruitworm, Sparganothis sulfureana, Tipworm	1.1 to 2.25 kg/ha	
	Fireworms	2.25 kg/ha	
Currants	Aphids, Currant borer, Fruitworm, Leaf roller, Mites,	1.1 to 1.25 kg/ha	
Blackber ries, Boysenbe rriesLog an berries Raspberr ies	Leafhoppers	1.1 kg/ha	Apply specified dosage per hectare to foliage using approximately 1600 L of water for good coverage. Repeat as necessary but not within 7 days of harvest.
	Aphids, Leaf miners, Leaf rollers, Obscure root weevil	1.75 kg/ha	



	Raspberry root borer (Crown)	2.25 to 4.5 kg/ha	For control of root weevil and borer prior to harvest apply specified dosage per hectare to lower portion of canes and to the soil beneath the plants using approximately 1600 L of water. Rates up to 2.25 kg/ha may be applied within 3 days of harvest. Rates above 2.25 kg/ha should be applied only before fruit set or after crop is harvested.
	Obscure root weevil	1.75 kg/ha	
Grapes	Flea beetle, Grape berry moth, Grape cane girdler, Ater ampelogypper, Leafhoppers, Leaf rollers, Mites	625 g to 1.25 kg/1000 L	Apply specified dosage in 1000 L of water as a full coverage spray.* Repeat as necessary but not more than 4 times per season. Minimum dosage specified may be applied up to harvest. Higher rates up to 875 g require a 5-day interval to harvest and above 875 g a 28-day interval to harvest.
	Red-banded leaf roller	1.25 to 2 kg/1000 L	
Melons Pumpkins	Spotted cucumber beetle, Striped cucumber beetle	1.1 kg/ha	Apply early in season when cucumber beetles first appear. A second application may be made if needed. A minimum 48-hour interval between each spray application and entry into the field must be observed. Do not apply within 14 days of harvest.

Strawberries	Aphids, Meadow spittlebug, Oblique-banded leaf roller, Strawberry leaf rollers, Omnivorous leaf tier, Obscure root weevil, Pea leaf weevil, Whitefly	1.1 kg/ha	Apply specified dosage in 1600 L of water per hectare as a full coverage spray.* Repeat as necessary. Do not apply within 5 days of harvest.
NUTS Walnuts	Aphids, Codling moth, European red mite, Filbertworm	925 g to 2.25 kg/1000 L	Apply specified dosage in 1000 L of water as a full coverage spray.* Repeat as necessary. Do not apply after husks split. Do not graze livestock in treated groves for 21 days after treatment.

VEGETABLES Beans (Green and Dry)	Leaf rollers, Mites	550 g to 1.1 kg/ha	Apply specified dosage per hectare in sufficient water to give complete coverage. Repeat as necessary. Do not apply within 3 days of harvest.
	Aphids, Green cloverworm, Leaf miners	850 g to 1.1 kg/ha	
	Leafhoppers, Mexican bean beetle, Spotted cucumber beetle, Striped cucumber beetle,	1.1 kg/ha	

Broccoli, Brussels Sprouts, Cabbage, Cauliflower	Aphids, Cabbage looper, Imported cabbageworm, Diamondback moth	1.1 to 1.75 kg/ha	Apply specified dosage per hectare. Repeat as necessary. Do not apply within 7 days of harvest for Brussels sprouts, 15 days of harvest for broccoli and cauliflower, nor within 21 days of harvest for cabbage.
	Cabbage maggot	1kg/1000 L	Mix specified dosage in 1000 L of water. Apply 115 to 170 mL per plant at or immediately after transplanting. Maintain sufficient agitation of the wetttable powder suspensions to avoid setting out in the spray tank.
Cucumbers	Spotted cucumber beetles, Striped cucumber beetles	1.1 kg/ha	Apply specified dosage per hectare in sufficient water to give complete coverage. Repeat as necessary. Do not apply within 6 days of harvest.
Potatoes	Colorado potato beetle,	550 to 850 g/ha	Apply specified dosage per hectare. Repeat as necessary. Do not apply within 7 days of harvest.
	European corn borer, Flea beetle, Leafhoppers, Spittlebug, Tarnished plant bug,	1.1 to 1.75 kg/ha	
	Aphids	1.75 kg/ha	
Spinach	Aphids, Leaf miners, Mites	850 g to 1.1 kg/ha	Apply specified dosage per hectare. Repeat as necessary. Do not apply within 14 days of harvest.
Tomatoes	Colorado potato beetle	850 g/ha	Apply specified dosage per hectare. Repeat as necessary. Do not treat within 2 days of harvest.

	Drosophila, Fruitworm, Green stink bug, Hornworm, Leaf miners, Whitefly	850 g to 1.1 kg/ha	
	Aphids, European corn borer, Flea beetles, Leafhoppers, Thrips	1.1 to 1.75 kg/ha	See previous column.
Turnips, Rutabagas, (Quebec & the Atlantic Provinces Only)	Cabbage maggot	37.5 g/100 m of row	AZINPHOS-M 50W is recommended for use as a postplanting soil drench. Mix specified dosage in 15 to 20 L of water and apply to 100 m of row in a 10 to 15 cm band sprayed over the row. Repeat as needed with a minimum of 30 days between applications. Do not apply within 35 days of harvest. Consult local authorities for proper timing of application.
ORNAMENTALS Ornamentals Nursery Plants, Forest & Shade Trees	Aphids, Euonymus scale, Lace bugs, Leafhoppers, Mites, Thrips	925 g to 1.25 kg/1000 L	Apply specified dosage per 1000 L of water. Spray all foliage surfaces including the underside of leaves for complete coverage.
	European elm scale	2 to 2.5 kg/1000 L	For control of brown soft scale and European scale use 50 g/10 L. Repeat as necessary.
	Brown soft scale	2.5 kg/ 1000 L	
	Cone midge, Cone moth	5 to 10 kg/1000 L	Apply specified dosage per 1000 L of water. Time applications to coincide with moth flight when cones are open for pollination. Thorough coverage of cones is necessary for maximum control. Repeat as necessary.

	Coneworm, Seedworm	3.75 kg/1000 L	
	European pine shoot moth	850 g to 1.75 kg/ha	Apply specified dosage per hectare in sufficient water for good coverage. Time applications to coincide with moth flights. For application to individual trees use 2 g/L.
	Nantucket pine tip moth	1.75 kg/ha	

\*This concentration is calculated for conventional sprayers. When lower volumes of spray are applied per hectare, as with low-pressure, low-volume, airplane, or mist-type equipment, increase the concentration of AZINPHOS-M 50W in the spray mixture in order to apply amount of AZINPHOS-M 50W per hectare equivalent to a full coverage spray.

#### PRECAUTIONS

Do not use on other crops used for food or forage. Treated orchards and groves should not be pastured within 21 days of treatment. Use only according to label directions. Application at rates above those shown may result in illegal crop residues. Do not use on food crops grown in the greenhouse. To protect fish and wildlife, do not contaminate streams, lakes, or ponds.

This product is highly toxic to bees exposed to direct treatment or residues on crops. Do not apply when crop is in bloom or allow spray to drift towards beehives. Protective information may be available from your local government extension specialist.

#### DANGER

Poisonous if swallowed, inhaled, or absorbed through the skin. Do not get in eyes or on skin. Wear protective clothing, natural rubber gloves, and goggles. Do not breathe dust or spray mist. Wear a mask or respirator effective for protection against AZINPHOS-M 50W. Information as to suitable types of masks or respirators is available from your dealer. Do not contaminate feed or food. KEEP OUT OF REACH OF CHILDREN. Keep all unprotected persons out of the operating area or vicinity where there may be danger of drift. Workers who must enter treated fields within 2 days of application should wear protective clothing. Wash hands, arms, and face thoroughly with soap and warm water before eating or smoking. Wash all contaminated clothing with soap and hot water before reuse.

#### FIRST AID

In case of poisoning call a physician immediately. Have patient lie

down and keep quiet. If swallowed, vomiting should be induced. Administer water freely and induce vomiting by giving one dose (15 mL) of syrup of ipecac. If vomiting does not occur within 10 to 20 minutes, administer second dose. If syrup of ipecac is not available, induce vomiting by sticking finger down throat. Repeat until vomit fluid is clear. The patient should be lying down with the head below the foot level and facing down or to one side. Professional medical assistance should be secured immediately. DO NOT INDUCE VOMITING TO AN UNCONSCIOUS PERSON OR TO PERSONS IN A CONVULSIVE STATE. If on skin, remove contaminated clothing and wash skin immediately with soap and warm water. If eyes are contaminated, wash immediately with flowing water for at least 15 minutes.

#### POISON TOXICOLOGICAL INFORMATION

**SYMPTOMS OF POISONING:** A sense of "tightness" in the chest. Sweating. Contracted pupils. Stomach pains. Vomiting and diarrhea.

**ANTIDOTE-** Administer atropine sulfate in large therapeutic doses. Repeat as necessary to the point of tolerance. 2-PAM is also antidotal and may be administered in conjunction with atropine.

Compound inhibits cholinesterase resulting in stimulation of the central nervous system, the parasympathetic nervous system, and the somatic motor nerves. Do not give morphine. Watch for pulmonary edema, which may develop in serious cases of poisoning even after 12 hours. At first sign of pulmonary edema, the patient should be placed in the oxygen tent and treated symptomatically.

#### CONTAINER DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

1. Completely empty the container into application equipment. Rinse the emptied container thoroughly and add the rinsings to the spray mixture in the tank.
2. Follow Provincial instruction for any additional cleaning of the container prior to its disposal.
3. Do not reuse container. Make the container unsuitable for further use.
4. Dispose of the empty container in accordance with Provincial requirements.
5. For information on the disposal of unused, unwanted product and the cleanup of spills, contact the regional office of Conservation and Protection, Environment Canada.

**NOTICE TO USER:** This control product is to be used only in accordance with the directions on this label. It is an offence under the Pest Control Products act to use a control product under unsafe conditions.

NOTE TO BUYER: Seller's guarantee shall be limited to the terms set out on the label and subject thereto, the buyer assumes the risk to persons or property arising from the use or handling of this product and accepts the product on that condition.

® AZINPHOS-M is a registered trademark of Chemagro Limited, UNITED AGRI PRODUCTS is a registered user.

94.02.09

SUPPLEMENTAL LABEL

FOR EUROPEAN APPLE SAWFLY ON APPLES

CLEAN CROP(R)

AZINPHOS-M(R) 50W

INSECTICIDE

RESTRICTED

FOR INSECT CONTROL ON FIELD, FRUIT, VEGETABLE AND ORNAMENTAL CROPS

GUARANTEE:

Azinphos-methyl ..... 50%

Store in cool dry place. Protect from heat.  
Keep away from open flame. Do not heat.

REGISTRATION NO. 22087 PEST CONTROL PRODUCTS ACT

DANGER POISON

KEEP OUT OF REACH OF CHILDREN

CALL A DOCTOR IN CASE OF ACCIDENT

READ THE LABEL BEFORE USING

NET: 2 kg

UNITED AGRI PRODUCTS  
R.R. #2

Dorchester, Ontario  
N0L 1G5

DIRECTIONS FOR USE

CROP	PEST	RATE	REMARKS
Apples	European Apple Sawfly	750 g/100 L	Apply specified dosage in 1000 L of water as a full coverage spray*. Do not apply more than once per season. Normal timing for this pest is at the calyx growth stage. Do not apply within 7 days of harvest. Consult your local agricultural authorities for specific directions on timing of application.

\*This concentration is calculated for conventional sprayers. When lower volumes of spray are applied per hectare, as with low-pressure, low - volume, airplane, or mist-type equipment, increase the concentration of AZINPHOS-M 50W in the spray mixture in order to apply amount of AZINPHOS-M 50W per hectare equivalent to a full coverage spray.

PRECAUTIONS

Do not use on other crops used for food or forage. Treated orchards and groves should not be pastured within 21 days of treatment. Use only according to label directions. Application at rates above those shown may result in illegal crop residues. Do not use on food crops grown in the greenhouse. To protect fish and wildlife, do not contaminate streams, lakes, or ponds.

This product is highly toxic to bees exposed to direct treatment or residues on crops. Do not apply when crop is in bloom or allow spray to drift towards beehives. Protective information may be available from your local government extension specialist.

DANGER

Poisonous if swallowed, inhaled, or absorbed through the skin. Do not get in eyes or on skin. Wear protective clothing, natural rubber gloves, and goggles.



Do not breathe dust or spray mist. Wear a mask or respirator effective for protection against AZINPHOS-M 50W. Information as to suitable types of masks or respirators is available from your dealer.

Do not contaminate feed or food. Keep out of reach of children. Keep all unprotected persons out of the operating area or vicinity where there may be danger of drift. Workers who must enter treated fields within 2 days of application should wear protective clothing. Wash hands, arms, and face thoroughly with soap and warm water before eating or smoking. Wash all contaminated clothing with soap and hot water before reuse.

#### FIRST AID

In case of poisoning call a physician immediately. Have patient lie down and keep quiet. If swallowed, vomiting should be induced. Administer water freely and induce vomiting by giving one dose (15 mL) of syrup of ipecac. If vomiting does not occur within 10 to 20 minutes, administer second dose. If syrup of ipecac is not available, induce vomiting by sticking finger down throat. Repeat until vomit fluid is clear. The patient should be lying down with the head below the foot level and facing down or to one side. Professional medical assistance should be secured immediately. DO NOT INDUCE VOMITING TO AN UNCONSCIOUS PERSON OR TO PERSONS IN A CONVULSIVE STATE.

If on skin, remove contaminated clothing and wash skin immediately with soap and warm water. If eyes are contaminated, wash immediately with flowing water for at least 15 minutes.

#### POISON TOXICOLOGICAL INFORMATION

SYMPTOMS OF POISONING: A sense of "tightness" in the chest. Sweating. Contracted pupils. Stomach pains. Vomiting and diarrhea.

ANTIDOTE- Administer atropine sulfate in large therapeutic doses. Repeat as necessary to the point of tolerance. 2-PAM is also antidotal and may be administered in conjunction with atropine.

Compound inhibits cholinesterase resulting in stimulation of the central nervous system, the parasympathetic nervous system, and the somatic motor nerves. Do not give morphine. Watch for pulmonary edema, which may develop in serious cases of poisoning even after 12 hours. At first sign of pulmonary edema, the patient should be placed in the oxygen tent and treated symptomatically.

#### CONTAINER DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

1. Completely empty the container into application equipment. Rinse

the emptied container thoroughly and add the rinsings to the spray mixture in the tank.

2. Follow Provincial instruction for any additional cleaning of the container prior to its disposal.
3. Do not reuse container. Make the container unsuitable for further use.
4. Dispose of the empty container in accordance with Provincial requirements.
5. For information on the disposal of unused, unwanted product and the cleanup of spills, contact the regional office of Environmental Protection, Environment Canada.

NOTICE TO USER: This control product is to be used only in accordance with the directions on this label. It is an offence under the Pest Control Products act to use a control product under unsafe conditions.

NOTE TO BUYER: Seller's guarantee shall be limited to the terms set out on the label and subject thereto, the buyer assumes the risk to persons or property arising from the use or handling of this product and accepts the product on that condition.

(R) AZINPHOS-M is a registered trademark of Chemagro Limited, UNITED AGRI PRODUCTS is a registered user.

\*\*\*\*\*

This label transcript service is offered by the Pest Management Regulatory Agency to provide efficient searching for label information. This service and this information do not replace the official hard-copy label. The PMRA does not provide any guarantee or assurance that the information obtained through this service is accurate, current or correct, and is therefore not liable for any loss resulting, directly or indirectly, from reliance upon this service.

+))