

85.12.31

MULTI-PURPOSE TETRALATE INSECTICIDE E.C. CODE 226.25

COMMERCIAL

COMBINING TWO SYNTHETIC PYRETHROID COMPOUNDS

A WATER EMULSIFIABLE INSECTICIDE CONCENTRATE

GUARANTEE :

TETRAMETHRIN .....	2.50%
RELATED COMPOUNDS .....	0.34%
RESMETHRIN .....	2.50%
RELATED COMPOUNDS .....	0.46%

\*CIS/TRANS ISOMER RATIO : MAX 30% (+) CIS AND  
MIN. 70% (+) TRANS.

REGISTRATION NO. 16126

PEST CONTROL PRODUCTS ACT

READ THE LABEL BEFORE USING

TETRALATE : REGISTERED TRADEMARK OF  
FAIRFIELD AMERICAN CORPORATION

FAIRFIELD CHEMICAL CANADA LTD  
238 WILSON AVENUE  
NEWARK, NJ 07105  
USA

#### DIRECTIONS FOR USE

For use in non-food areas such as dwellings, offices, hospitals, hotels, motels and schools.

This concentrate should be diluted with water before use, at a rate of 1 part concentrate to 9 parts water. The diluted spray should be applied in conventional mechanical or compressed air spraying or fogging equipment. It is not recommended for use in thermal generating equipment.

To control ants, centipedes, cockroaches, earwigs, crickets, silverfish, sowbugs and spiders: Apply as a coarse droplet spray directed into cracks, crevices and other hiding places along baseboards, around the base of machinery and under pallets. Treat all returned bottle cartons and cases. Spray thoroughly contacting insects directly whenever possible. Spray badly infested areas daily. In other areas, repeat treatments twice weekly or when reinfestation occurs.

To control flies, gnats, mosquitoes and small flying moths: Close doors and windows and shut off ventilating systems. Apply as a fog or fine mist directing the spray toward the upper portions of the room and towards windows and other light sources which attract these insects. Thoroughly fill the area with mist using 50 to 100 mL per 100 cubic meters of space. Keep area closed for at least 10 minutes. Vacate after treatment and ventilate before reoccupying. Repeat treatment as needed.

On ornamental Plants: This concentrate should be diluted with water before use, at a rate of 1 part concentrate to 79 parts water (12 mL per liter or 1/2 liter per 40 liters). The diluted spray should be applied in conventional mechanical or compressed air spraying or fogging equipment. It is not recommended for use in thermal generating equipment.

To control aphids, two-spotted spider mites and white flies (adults) on ornamental house and garden plants as listed :

African Violets	Gloxinias
Azaleas	Ivies
Christmas Cherries	Lantanas
Chrysanthemums	Neanth Bella Palms
Cinerarias	Petunias
Diffenbachias	Roses
Fuschias	Snapdragons

Outdoor Plants :

Azaleas	Petunias
Chrysanthemums	Roses
Euonymus	Snapdragons
Ivies	Zinnias
Marigolds	

Apply as a fog or fine mist, spraying plants with a sweeping motion so that the mist drifts into the plants. Direct spray so that the upper and lower portions of leaves are contacted. Use caution on new growth and tender foliage. When using outdoors, spray when air is still. Repeat as necessary.

To control cankerworms, garden webworms, tent caterpillars and the above pests on ornamental crabapple, cherries and plum trees, spray as above. Do not spray on bearing fruit trees.

PRECAUTION: KEEP OUT OF REACH OF CHILDREN. Harmful if swallowed. Exposed foods should be removed or covered during treatment. All food preparation surfaces should be covered during treatment or thoroughly cleaned before using. Wash all contaminated clothing with soap and hot water before reuse. Remove pets, birds and cover fish aquariums before spraying.

\*\*\*\*\*

This label transcript service is offered by the Pest Management Regulatory Agency to provide efficient searching for label information. This service and this information do not replace the official hard-copy label. The PMRA does not provide any guarantee or assurance that the information obtained through this service is accurate, current or correct, and is therefore not liable for any loss resulting, directly or indirectly, from reliance upon this service.

+))