

89.12.22

CHEMCO LV 96

Emulsifiable Concentrate

- Low Volatile Farm and Industrial Weed Killer
- Contains 2,4-D
- For air or ground selective spraying of wheat, barley, rye, oats, corn, pastures, turf, for the control of certain annual and perennial weeds and for woody plant growth.

AGRICULTURAL

WARNING

POISON

READ THE LABEL BEFORE USING. KEEP OUT OF REACH OF CHILDREN.

GUARANTEE: 2,4-D equivalent.....564 g/L  
(as low volatile esters of 2,4-D)

REGISTRATION NO. 21265  
PEST CONTROL PRODUCTS ACT

NET CONTENTS:

PRECAUTIONS: Causes irritation of skin and eyes. Do not get in eyes. Avoid contact with skin and clothing. Remove and wash contaminated clothing and shoes before re-use.

FIRST AID: If swallowed, do not induce vomiting. GET MEDICAL ATTENTION. In case of eye contact, flush eyes with plenty of water for at least 15 minutes and get medical attention. If on skin, wash with soap and plenty of water.

TOXICOLOGICAL INFORMATION: Treat Symptomatically.

NOTICE TO USER: This control product is to be used only in accordance with the directions on this label. It is an offence under the PEST CONTROL PRODUCTS ACT to use a control product under unsafe conditions.

NOTICE TO BUYER: Seller's guarantee shall be limited to the terms set out on the label and subject thereto, the buyer assumes the risk to persons or property arising from the use or handling of this product and accepts the product on that condition.

AGRICULTURAL CHEMICAL: Do Not Ship Or Store With Food, Feeds, Drugs or Clothing

DIVISION OF:  
OLIVER INDUSTRIAL SUPPLY LTD.  
236-36 ST. NORTH  
LETHBRIDGE, ALBERTA  
T1J 4B2

\* Trademark of DowElanco

## DIRECTIONS FOR USE

### GENERAL INSTRUCTIONS - CROP USE

**GROUND APPLICATION:** Apply 50-200 L of spray solution per hectare depending on the type of application equipment used. Use sufficient water for even distribution. Spray at low pressures (200-275 kPa) when the weeds are actively growing.

**AERIAL APPLICATION:** (including fixed and rotary wing aircraft). Apply a minimum of 30 L of spray solution per hectare depending on the type of application system being used. Use boom pressures of 235 kPa or less. Avoid placing nozzles where spray will enter wing tip vortices.

**To Prepare A Spray:** Add half the required amount of water to the spray tank, then the OCHEMCO LV 96 with agitation, and finally the balance of the water with continued agitation.

**WARNING:** If preparing straight oil mixtures, do not let water get into the product or the finished mixture. Note: This product in water forms an emulsion, not a solution, which tends to separate on standing. Agitate to prevent such separation and ensure uniformity of spray mixture.

**Spot Treatment:** For knapsack application of OCHEMCO LV 96 for spot treatment of weeds such as thistles, mix 200 mL of product in 10 L of water. Wet all foliage thoroughly.

**Selective Weed Control in Small Grains:** Weeds differ in their susceptibility to OCHEMCO LV 96 and not all types can be controlled satisfactorily in crops. The amount of OCHEMCO LV 96 to use will depend upon the susceptibility and whether the crops will tolerate this amount. See chart.

**Wheat, Barley, Rye:** Spray from the 3-leaf expanded stage (15 cm tall) to just before the flag-leaf (shot-blade) stage. Sprays may be used after the hard dough stage to control weeds which might interfere with harvesting. To avoid crop injury, do not treat during boot and flowering stages. Winter wheats and fall rye should be treated in early spring, as soon as weeds appear and before the crop reaches the shot-blade stage.

**Oats:** Oats are more sensitive to 2,4-D than wheat or barley,

especially when 15 cm tall (3-to-6 leaf stage). If possible, treat oats from emergence to the 3-leaf stage or from the 6-leaf to the flag-leaf stage.

#### APPLICATION

Suggested Amounts of OCHEMCO LV 96 for Weed Control in Cereals (not underseeded to legumes)

Mixed Weedy Growth	Stage of Weed Kind of Weather	L/ha	Hectares Treated/ 20 L
Susceptible weeds such as: mustard (except dog and green tansy), flixweed, bluebur, burdock, cocklebur, *field horsetail, stinkweed, goatsbeard, prickly lettuce, *hoary cress, kochia, lambsquarters, plantain, ragweeds, Russian knapweed, Russian thistle, shepherd's purse, annual sunflower, sweet clover, vetch, wild radish.	Seedling (2-4 leaves) 600 mL 33 Growing rapidly	600 mL	34
	Weeds in bud Dry, cool weather Heavy infestation	900 mL	22
Hard-to-kill weeds such as: biennial wormwood, docks, dandelion, dog mustard, field peppergrass, gumweed, hairy galinsoga, hedge bindweed, ladythumb, oak leaf goosefoot, redroot pigweed, Russian thistle, smartweed, annual sow-thistle, tansy, tartary buckwheat, tumble-weed, *blue lettuce, *Canada thistle, *field bindweed, *leafy spurge, *perennial sowthistle, ** wild buckwheat	Seedling (2-4 leaves) Growing rapidly	1.0+	20
	Weeds in bud Dry or cool weather Heavy infestation	1.5+	13

+ The higher rates may cause deformities and delayed maturity which will be offset by a higher relative yield from a less weedy crop.

\* Top growth control only.

\*\* Better control will result from 2 treatments applied one week apart at 0.6 L/ha 2,4-D

Corn: Post-Emergence - Treat with 700 mL/ha when corn plants are 15 to 20 cm tall and most weeds have germinated but are still in seeding stage. Use drop nozzles to keep spray off corn foliage. Do not cultivate until at least 2 weeks after treatment.

Established Grasses: Eastern Canada - For the control of yellow rocket and other susceptible winter annual weeds, spray with 1.7 L of OCHEMCO LV 96/ha Application should be made in early fall.

Established Alfalfa and Grasses: Western Canada - To control susceptible perennials or winter annual, spray 800 mL to 1.5 L/ha before growth of legumes and grasses starts in the spring.

Rangeland, Grass Pasture and Turf: To control susceptible weeds, use 1.5 to 2.75 L/ha in enough water for coverage.

WARNING: Do not use on lawns of creeping grasses such as bent, except for spot spraying, or on freshly seeded turf until grass has become well established. Most legumes are usually damaged or killed.

#### GENERAL INSTRUCTIONS - NON-CROP USE

GROUND APPLICATION: Apply 50-3000 L of spray solution per hectare depending on the type of application system being used. Use sufficient water to ensure uniform coverage.

AERIAL APPLICATION: (including fixed and rotary wing aircraft). Apply a minimum of 30 L of spray solution per hectare depending on the type of application system being used. Use boom pressures of 235 kPa or less. Avoid placing nozzles where spray will enter wing tip vortices.

ROADSIDE AND GENERAL WEED CONTROL: For taller weed growth such as in vacant lots, on roadsides and along fence rows, sufficient spray solution should be used to wet all foliage thoroughly. Use ESTERON LV- 600 at the rate of 3.75 L per hectare. Deeprooted perennial weeds such as bindweed and Canada thistle, may require repeated applications as new growth appears.

WOODY PLANTS: To control 2,4-D susceptible species, spray brush up to 2.5 m tall after foliage is well developed. When using ground application equipment, mix 8.0 L of OCHEMCO LV 96 in 1000 L of water and spray to wet all foliage, shoots stems and bark. With good growing conditions and high soil moisture, sprays may be used up to 2 or 3 weeks before normal frost time, but

spraying soon after leaves are fully open in the spring usually gives best results. Control may be less effective during hot, dry weather when deep soil moisture is deficient. For 2,4-D resistant species use TORDON\* 101 Mixture Herbicide.

When using aerial application equipment apply 6.75 L of OCHEMCO LV 96 in a minimum of 30 L of spray solution per hectare.

#### USE PRECAUTIONS

##### WARNING

FOR AERIAL APPLICATIONS: Do not spray when winds exceed 8 kmph. Do not spray in dead clam near sensitive plants. The "cloud" of suspended droplets may drift onto sensitive plants when the wind comes up. Spray only when wind is blowing away from a sensitive crop, shelterbelt or garden.

SENSITIVE PLANTS: Vegetables, flowers, grapes, fruit trees and other desirable plants are sensitive to 2,4-D even in minute quantities. Care should be taken to avoid spraying these types of plants or allowing spray mist to drift onto these plants during both their growing and dormant periods. Coarse sprays are less likely to drift. At higher temperatures, vapourization may cause injury to susceptible plants growing nearby.

This product may cause damage to lawns or pastures if applied before the grass is well established. In addition, most legumes may be damaged or killed. Allow 24 hours between application of product and grazing/cutting for hay.

STORAGE: This product must be stored away from fertilizers, seeds, insecticides, fungicides or other herbicides intended for use on 2,4-D sensitive crops.

USE: When the container is empty triple rinse and add the rinse water to the spray tank. Care must be taken during application to avoid contaminating irrigation ditches or domestic water supplies.

SPRAYER CLEANUP AND DISPOSAL OF WASTE: When spraying is complete, rinse all equipment and the original container with water and dispose of rinse water by burying in noncropland away from water supplies. Avoid usingsprayers which contained ,4-D to apply other pesticides to sensitive crops.

##### CONTAINER DISPOSAL:

1. Rinse the emptied container thoroughly and add the rinsings to the spray mixture in the tank.
2. Follow provincial instructions for any required additional cleaning of the container prior to its disposal.
3. Make the empty container unsuitable for further use.
4. Dispose of the container in accordance with provincial requirements.

5. For information on the disposal of unused, unwanted product and the cleanup of spills contact the regional office of Conservation and Protection, Environment Canada.

NOTE: Local conditions may affect the use of herbicides. Provincial agricultural authorities issue recommendations to fit local conditions. Be sure that use of this product conforms to all applicable regulations.

If this product is exposed to temperatures below -20'C, it should be warmed to at least 5'C and mixed thoroughly before using.

\*\*\*\*\*

This label transcript service is offered by the Pest Management Regulatory Agency to provide efficient searching for label information. This service and this information do not replace the official hard-copy label. The PMRA does not provide any guarantee or assurance that the information obtained through this service is accurate, current or correct, and is therefore not liable for any loss resulting, directly or indirectly, from reliance upon this service.

+))