

22-NOV-1999

dispar

TRAP-N-A-SAK (T.M.)

RODENTICIDE

THE SWEET CORN TRAP RODENTS CAN'T RESIST

KILLS RATS AND MICE

COMMERCIAL

5.45 kg BULK BUCKET

WITH DIPHACINONE

A PELLETIZED RODENTICIDE
"CONTAINING SWEET CORN"
READY MIXED, USE INDOORS
AND OUTDOORS.

GUARANTEE:

DIPHACINONE..... 0.005%

REGISTRATION NO. 21436 PEST CONTROL PRODUCTS ACT

KEEP OUT OF REACH OF CHILDREN

READ LABEL BEFORE USING

REGISTERED TRADEMARK

DISPAR,
DIVISION OF
VETOQUINOL N.-A. INC.
LAVALTRIE, QC. CANADA
J0K 1H0

INDICATIONS: For control of Norway rats (*Rattus norvegicus*), roof rat (*Rattus rattus*) and house mice (*Mus musculus*) in and around homes, industrial and agricultural buildings, and similar man-made structures. Treated baits must be placed in tamper-proof bait boxes or in location not accessible to children, pets, domestic animals or wildlife. Do not place bait

in areas where there is a possibility of contaminating food or surfaces that come in contact with food.

SELECTION OF TREATMENT AREAS: Determine areas where rats and/or mice will most likely find and consume the bait. Generally, these areas are along walls, by gnawed openings, in or beside burrows, in corners and concealed places, between floors and walls, or in locations where rodents or their signs have been observed. Protect bait from rain or snow. Remove as much food as possible.

DIRECTIONS: RATS: Apply 100 to 500 g of bait per placement. Maintain an uninterrupted supply of fresh bait for at least 10 days.

MICE: Apply 7 to 15 g (1-2 tablespoons) of bait at 1-3 m intervals. Larger placements (up to 60 g) may be needed at points of very high mouse activity. Maintain an uninterrupted supply of fresh bait for at least 15 days.

RATS AND MICE: Replace contaminated or spoiled bait immediately. Collect and dispose of all dead animals and unconsumed bait according to local requirements. To prevent reinfestation, eliminate food, water and harbourage as much as possible. If reinfestation does occur, repeat treatment. Where continuous source of infestation is present, establish permanent bait stations and replenish bait as needed.

PRECAUTIONS: KEEP OUT OF THE REACH OF CHILDREN. Keep away from humans, domestic animals and pets. If swallowed, this material may reduce the clotting ability of the blood and cause bleeding.

FIRST AID: If swallowed, give patient one or two glasses of water and induce vomiting by inserting finger down throat; repeat until vomit fluid is clear. Call a physician.

TOXICOLOGICAL INFORMATION: Vitamin K1 in the form of intravenous, intramuscular or subcutaneous injections, or by oral ingestion are suggested remedial treatments for anticoagulant poisoning. The severity of the case, measured by establishing prolonged prothrombin time (P.T.), will determine appropriate therapy. Monitoring P.T. will indicate the necessity of repeated treatments.

DISPOSAL: Do not reuse the container. Dispose of the container in accordance with provincial requirements. For information on the disposal of unused, unwanted product and the cleanup of spills, contact the regional office of Conservation and Protection, Environment Canada.

LIMITATION OF WARRANTY: Seller's guarantee shall be limited to the terms set out on the label and, subject thereto, the buyer assumes the risk to persons or property arising from the use or handling of this product and accepts the product on that condition.

NOTICE TO USER: This control product is to be used in accordance with the directions on this label. It is an offense under the Pest Control Products Act to use a control product under unsafe conditions.

This label transcript service is offered by the Pest Management Regulatory Agency to provide efficient searching for label information. This service and this information do not replace the official hard-copy label. The PMRA does not provide any guarantee or assurance that the information obtained through this service is accurate, current or correct, and is therefore not liable for any loss resulting, directly or indirectly, from reliance upon this service.

+))