

93.10.29

apa

LOUSE KILL

POUR-ON EMULSIFIABLE LIQUID

For control of cattle and hog lice
Aids in the control of Hog Sarcoptic Mange

AGRICULTURAL

GUARANTEE:

PHOSMET..... 11.6%

REGISTRATION NO. 23122
PEST CONTROL PRODUCTS ACT

READ THE LABEL BEFORE USING

PROTECT FROM FREEZING

DANGER POISON

WARNING:

KEEP OUT OF REACH OF CHILDREN

May be harmful if swallowed, inhaled or absorbed through the skin. Do not get in eyes, on skin or on clothing. Do not breathe in mist. Wear rubber gloves, goggles and protective clothing. In case of skin contact, wash immediately with soap and water. Wash all contaminated clothing with soap and hot water before re-use. Do not store near heat or open flame.

See toxicological information.

A.P.A.
division of sanofi sante animale
canada, inc.
Victoriaville, Qc, G6P 1B1, Canada

NET CONTENTS: 1 L

code 21250

DIRECTIONS FOR USE

POUR-ON METHOD: Dilute one volume of Louse Kill with five volumes of water and stir vigorously.

One volume of Louse Kill makes 6 volumes of pour on solution.
6 litres will treat:

Animal Weight (kg)	100	150	200	250	300	350	400 or over
Number of Animals treated	100	66	50	40	33	28	25

Cattle Lice: Apply 60 mL of the freshly diluted mixture per 100 kg of body weight (to a maximum of 230 mL per head) down the centre line of the animal's back. Cattle may become reinfested 3 weeks after treatment.

Hog Lice: Apply 60 mL of the freshly diluted mixture per 100 kg of body weight down the centre line of the animal's back. Re-treat as necessary but not more frequently than every 7 days.

Hog Mange: Apply 60 mL of the freshly diluted mixture per 100 kg of body weight down the centre line of the animal's back. A second application 14 days later may be necessary.

SPRAY METHOD: Dilute one volume of Louse Kill with 100 volumes of water and stir vigorously.

Cattle Lice: Apply the fresh mixture as a high-pressure spray directly to wet the skin (not just the hair) of each animal to "run-off". Use about 3.5 L of the diluted spray per adult animal. Lesser amounts will permit run-off for younger animals. Spray entire body, giving particular attention to lice infested areas. Cattle may become re-infested 5 weeks after treatment.

Hornflies: Apply the fresh mixture as a high-pressure spray directly to wet the skin (not just the hair) of each animal to "run-off". Use about 3.5 L of the diluted spray per adult animal. Lesser amounts will permit run-off for younger animals. Re-treat as necessary but not more frequently than every 7 days.

Hog Lice: Apply the fresh mixture as a high-pressure spray directly to wet the skin (not just the hair) of each animal to "run-off". Use about 1 L of the diluted spray per adult animal.

Lesser amounts will permit run-off for younger animals. Re-treat as necessary but not more frequently than every 7 days.

Notes: Do not use on lactating cattle. Do not apply within 7 days of slaughter. Do not use this product on cattle which are likely to be infested with grubs as host-parasite reaction may occur. Symptoms of host-parasite action usually appear 24 to 96 hours after treatment and include staggering, or rarely posterior paralysis, salivation, or bloat. Consult your veterinarian should these symptoms occur. Atropine is contraindicated. Solvent irritation may occur in sensitive skinned animals, symptoms are increased activity, a repeated getting up and down, tail swishing, and a twitching of the skin. Symptoms occur within 5 to 15 minutes after treatment and subside within one hour without complications. Atropine is contraindicated.

Dandruff may appear on treated areas which may be a blemish in show animals.

FIRST AID:

1. Keep airways open. Use artificial respiration if necessary.
2. If ingested and patient is conscious, induce vomiting. Do not induce vomiting if patient is unconscious or in a convulsive state.
3. Call physician.
4. Wash skin of material and vomitus.
5. Save suspected agent and vomitus if container is available, for disposal by physician.
6. In case of contact with skin, wash immediately with soap and water. If in eyes, flush with water.

TOXICOLOGICAL INFORMATION: Phosmet is an organophosphorous insecticide. Atropine is antidotal for poisoning in man and animals. Usual symptoms of organophosphorous poisoning in man include: headache, blurred vision, weakness, nausea, discomfort in the chest, vomiting, abdominal cramps, diarrhea, salivation, sweating, pinpoint pupils. Symptoms of organophosphorous poisoning in animals usually occur within 30 to 60 minutes after application. These may include salivation, pinpoint pupils, laboured breathing, stiffness, muscle tremors, ataxia, and mild bloating.

PRECAUTIONS: Warning: Phosmet is a cholinesterase inhibitor, do not use this product on animals simultaneously or within seven days before or after treatment with, or exposure to cholinesterase inhibiting drugs, pesticides or chemicals. Do not treat sick, stressed or debilitated animals. Do not treat calves under 3 months of age. Consult you veterinarian for treatment of sick animals.

Do not apply with 7 days of slaughter. Do not retreat within 7 days. For use on beef cattle only. Do not treat dairy animals. Do not contaminate feed or foodstuffs. Avoid spray drift to adjoining food and forage crops. Toxic to fish. Keep out of streams, lakes, ponds and other fish-bearing waters.

STORAGE AND DISPOSAL:

STORAGE: Store in a cool, dry and ventilated area away from humans, pets, foodstuffs and feed. Do not store near hear or open flame. Protect from freezing.

DISPOSAL: Rinse the emptied container thoroughly and add the rinsings to application mixture. Follow Provincial instructions for any required additional cleaning of the container prior to its disposal. Make the empty container unsuitable for further use and dispose in accordance with Provincial requirements. For information on the disposal of unused, unwanted product and the cleanup of spills, contact the regional office of Environmental Protection, Environment Canada.

NOTICE TO BUYER: Seller's guarantee shall be limited to the terms set out on the label and subject thereto, the buyer assume the risk to persons or property arising from the use or handling of this product and accepts the product on that condition.

NOTICE TO USER: This control product is to be used only in accordance with the directions on this label. It is an offense under the Pest Control Products Act to use a control product under unsafe conditions.

INTEG (TM) is protected by the following Patents: U.S.A. 4744161, 4680080, 4592572, 4675062, Canada 422793, and others

This label transcript service is offered by the Pest Management Regulatory Agency to provide efficient searching for label information. This service and this information do not replace the official hard-copy label. The PMRA does not provide any guarantee or assurance that the information obtained through this service is accurate, current or correct, and is therefore not liable for any loss resulting, directly or indirectly, from reliance upon this service.

+))