

97.12.03

APRON(R) COMBI FS FUNGICIDE

FOR TREATMENT OF SEED DESTINED  
FOR EXPORT ONLY

THIS IS A READY-TO-USE FORMULATION  
- DO NOT DILUTE

AGRICULTURAL

NET CONTENTS: 20 litres

Controls: Ascochyta, downy mildew and seedling "damping-off"  
diseases in all varieties of peas

GUARANTEE:

metalaxyl	140 g/L
thiabendazole	72 g/L
thiram	60 g/L

WARNING - POISON  
WARNING - EYE IRRITANT  
POTENTIAL SKIN SENSITIZER

READ THE LABEL BEFORE USING

KEEP OUT OF THE REACH OF CHILDREN

REGISTRATION NO.: 24288  
PEST CONTROL PRODUCTS ACT

Ciba-Geigy Canada Ltd.  
Crop Protection Division  
6860 Century Avenue  
Mississauga, Ontario  
L5N 2W5

SPECIAL USE RESTRICTION: This product can only be applied to pea  
seed which will be exported.

NOTICE TO USER: This control product is to be used only in  
accordance with the directions on this label. It is an offence  
under the PEST CONTROL PRODUCTS ACT to use a control product

under unsafe conditions.

NOTICE TO BUYER: Seller's guarantee shall be limited to the terms set out on the label and subject thereto, the buyer assumes the risk to persons or property arising from the use or handling of this product and accepts the product on that condition.

FIRST AID: In case of poisoning, call a physician immediately or contact a poison control centre. If on skin, remove contaminated clothing and wash skin thoroughly with soap and water. If in eyes, flush with clean water for 15 minutes, and obtain medical attention or contact a poison control centre. If swallowed, give the patient 1-2 glasses of water and cause vomiting by giving one dose (15 mL) of syrup of ipecac. If the patient does not vomit within 20 minutes, give a second dose. If syrup of ipecac is not available, give the patient 1-2 glasses of water and cause vomiting by inserting finger down the throat. Repeat with water until vomit fluid is clear. The patient should be lying down with the head below the level of the feet. Do not try to cause vomiting if the patient is unconscious or in a convulsive state.

PRECAUTIONS: KEEP OUT OF REACH OF CHILDREN.

May be harmful if swallowed. Avoid inhalation of dust. Wash hands and face after handling. Avoid contamination of feed and foodstuffs. Treat seed in a well ventilated area. Wear coveralls over normal work clothing and chemical resistant gloves when handling APRON COMBI FS, contaminated equipment or APRON COMBI FS treated seed. Do not consume alcoholic beverages immediately before or within 24 hours of working with APRON COMBI FS.

DECONTAMINATION AND DISPOSAL:

1. In cleaning of equipment or disposing of wastes, do not contaminate water used for human or animal consumption or by wildlife and aquatic life or for irrigation purposes.
2. Empty container thoroughly into the mixing tank.
3. Rinse the emptied container thoroughly and dispose of the rinsings in accordance with provincial requirements.
4. Follow provincial instructions for any required additional cleaning of the container prior to its disposal.
5. Make the container unsuitable for further use.
6. Dispose of the container in accordance with provincial requirements.
7. For information on the disposal of unused, unwanted product,

and the cleanup of spills, contact the Provincial Regulatory Agency or the manufacturer.

GENERAL INFORMATION: APRON COMBI FS is a ready to-use flowable formulation containing metalaxyl, thiabendazole, and thiram for protection against Ascochyta, downy mildew and seedling "damping-off" diseases in peas. Do not dilute with water. SHAKE WELL BEFORE USING.

#### DIRECTIONS FOR USE:

##### Dose

5 litres APRON COMBI FS per tonne (1000 KG) of seed.  
(One 20 litre jug will treat 4 tonnes [4000 KG] of seed).

##### Method of Application

APRON COMBI FS should be applied through continuous flow seed treaters. Adequate seed coverage is obtained if APRON COMBI FS is added at the start of the conveyer or mixing process.

It is essential that any continuous flow seed treatment equipment be calibrated for APRON COMBI FS before commercial use.

##### Volume

APRON COMBI FS is a ready-to-use formulation which should NOT be diluted further. It should be applied to seed using a volume of 500 mL of APRON COMBI FS per 100 kg of seed (5 L/tonne)

##### Storage after Treatment

Ensure that the seeds are of a satisfactory moisture content before storage; keep in a dry place, correctly labelled. Treated seed should be coloured and labelled: "FOR EXPORT ONLY. This seed has been treated with metalaxyl, thiabendazole and thiram fungicides. Poisonous to man and animals. Do not use for feed or food purposes. Store away from feeds and foodstuffs".

(R)APRON is a registered trademark of Ciba-Geigy Canada Ltd.

\*\*\*\*\*

This label transcript service is offered by the Pest Management Regulatory Agency to provide efficient searching for label information. This service and this information do not replace the official hard-copy label. The PMRA does not provide any

guarantee or assurance that the information obtained through this service is accurate, current or correct, and is therefore not liable for any loss resulting, directly or indirectly, from reliance upon this service.

+))