

2000/2001

ASSASSIN

KILLS RATS AND MICE

RAT & MOUSE BAIT PELLETS

REGISTRATION NUMBER 23165  
PEST CONTROL PRODUCTS ACT

DOMESTIC

GUARANTEE:

Diphacinone.....0.005%

READ THE LABEL BEFORE USING

KEEP OUT OF REACH OF CHILDREN

NET CONTENTS: 85 g (3oz.)  
20 x 85g (3 oz.)

Distributed By:  
CITADEL  
ANIMAL HEALTH  
CAMBRIDGE, ON N3C 2W4

**PRECAUTIONS:** KEEP OUT OF REACH OF CHILDREN, PETS, DOMESTIC ANIMALS and WILDLIFE. Do not expose treated bait in areas accessible to children, dogs, domestic animals and wildlife, except when enclosed within a tamper-resistant bait box. Keep away from feed or food stuffs. Do not place bait in areas where there is a possibility of contaminating food or surfaces that come in direct contact with food. Do not get in eyes, on skin or on clothing. Wash arms, hands and face with soap and water after handling and before eating or smoking. Keep out of lakes, streams or ponds.

**FIRST AID:** If in eyes, flush with water for 15 minutes. If on skin, wash with soap and water. If swallowed, give patient one or two glasses of water and induce vomiting by touching back of throat with a finger or a blunt object; repeat until vomit fluid is clear. Do not give anything by mouth to an unconscious person. Contact a physician or Poison Control Centre immediately.

**TOXICOLOGICAL INFORMATION:** If swallowed this material may reduce

the clotting ability of the blood and cause bleeding. If ingested, Vitamin K1, in the form of intravenous, intramuscular or subcutaneous injections, or by oral ingestion are suggested remedial treatments for anticoagulant poisoning. The severity of the case measured by establishing prolonged prothrombin times(P.T.) will determine appropriate therapy. Monitoring P.T. will indicate the necessity of repeated treatments.

**APPLICATION DIRECTIONS:**

**RATS:** Place 100 to 500 grams of bait(usually at intervals of 4.5 to 9 meters) per placement. Maintain an uninterrupted supply of fresh bait for 10 days or until signs of rat activity cease.

**HOUSE MICE:** Place 7 to 14 grams of bait at intervals of 2.5 to 3.5 meters per placement. Larger placements(up to 100 grams) may be needed at points of high mouse activity. Maintain an uninterrupted supply of fresh bait for 15 days or until signs of mouse activity cease.

**RATS/MICE:** Replace contaminated or spoiled bait immediately. Collect and dispose of unconsumed bait and dead exposed animals according to local or provincial requirements. To prevent reinfestation, eliminate food, water, and harborage as much as possible. If reinfestation does occur, repeat treatment. Where a continuous source of infestation is present, establish permanent bait stations and replenish bait as needed.

**DIRECTIONS FOR USE:** Use bait to control the Norway rat,(Rattus norvegicus), Roof rat(Rattus rattus) and House mouse(Mus musculus) in and around dwellings.

**SELECTION OF TREATMENT AREAS:** Determine areas where rats and/or mice will most likely find and consume the bait. Generally, these areas are along walls, by gnawed openings, in or beside burrows, in corners and concealed places, between floors and walls, or in locations where rodents or their signs have been observed. Remove as much food as possible.

**DISPOSAL:** Do not reuse empty container. Dispose of empty container by placing in household garbage.

**Rodent Disposal:** Dispose of dead rodents in garbage or by burying.

\*\*\*\*\*

This label transcript service is offered by the Pest Management Regulatory Agency to provide efficient searching for label

information. This service and this information do not replace the official hard-copy label. The PMRA does not provide any guarantee or assurance that the information obtained through this service is accurate, current or correct, and is therefore not liable for any loss resulting, directly or indirectly, from reliance upon this service.

+))