

91.10.02

((LABEL))

AGROCORP

TRIFLUREX 40 EC

Contains Trifluralin

11.3L

AGRICULTURAL HERBICIDE

DIRECTIONS FOR PRAIRIE PROVINCES

A Selective Herbicide for Pre-emergence Weed Control
in many Field Crops, Vegetables and Shelterbelts.

READ THE LABEL AND ATTACHED INSTRUCTIONS BEFORE USING

GUARANTEE:

TRIFLURALIN.....400g/L

REGISTRATION NO. 21816
PEST CONTROL PRODUCTS ACT

CAUTION

POISON

MANUFACTURED FOR:
AGRACORP INC.
BRANDON, MANITOBA

Manufactured In Canada by:
SHAMROCK AGRO-TECH INC.
607-4th Avenue East
Regina, Saskatchewan
S4N 4Z8

595 Exeter Road
London, Ontario
N6E 2Z2

STORAGE

If stored below -18 degrees C bring the contents to 15 degrees C for 24 hours
and shake well before using.

DISPOSAL

Rinse the emptied container thoroughly and add the rinsings to the spray mixture in the tank. Follow provincial instructions for any required additional cleaning of the container prior to its disposal. Make the container unsuitable for further use. Dispose of the container in accordance with provincial requirements. For information on disposal of unused, unwanted product and the clean-up of spills, contact the regional office of Conservation and Protection, Environment Canada.

NOTICE TO USER

This control product is to be used only in accordance with the directions on this label. It is an offence under the Pest Control Products Act to use a control product under unsafe conditions.

NOTICE TO BUYER

Seller's guarantee shall be limited to the terms set out on the label and subject thereto: the buyer assumes the risk to persons or property arising from the use or handling of this product and accepts the product on that condition.

Triflurex is a registered trademark of Makhteshim-Agan (America) Inc. Manufactured by Shamrock Agro-Tech Inc., Regina, Saskatchewan & London, Ontario.

INSTRUCTIONS FOR USE

Carefully read these instructions fully before using TRIFLUREX. Usage not in accordance with these instructions may result in inadequate weed control and may damage crop.

TRIFLUREX is a pre-emergence herbicide which is sprayed on and then mechanically mixed or incorporated into the soil. It kills weeds as they germinate. It does not kill established weeds. TRIFLUREX gives long lasting control of seedlings of annual grasses and broadleaf weeds detailed on the enclosed pamphlet.

WARNING

Special attention must be paid to the following aspects of usage:

- Rates must be accurately applied.
- Apply only on soil types indicated (this excludes peat and muck soils).
- Apply only to soils with less than 15% organic matter which are dry and in good working condition. Soils with the potential of becoming waterlogged must not be treated.
- Incorporation is an essential part of application. When correctly applied to crops under normal conditions, TRIFLUREX will not harm the crop. Any stress factor such as wet soil, incorrect planting depth, seedling disease, low temperatures, excessive salt in the soil or drought, could bring about damage to the crop from TRIFLUREX.
- It is recommended that the following crops not follow TRIFLUREX in the rotation cycle: Oats, Small seeded annual grasses such as Canary Seed

grass; Sugarbeets; Creeping Red Fescue; Timothy grass. Under normal conditions there will not be a carryover. However, the above recommendation is precautionary.

CAUTION

TRIFLUREX may affect crops adversely if not used strictly in accordance with these instructions. It can only be used on crops mentioned on this label. The only grain crops for which it is suitable for use are wheat and barley. It may not be used on other grain crops such as oats, rye or grass crops. Conditions causing seedling stress, such as wet soils, incorrect planting depth, seedling disease, low temperatures, excessive salt in the soil or drought, could bring about damage to the crop from TRIFLUREX.

Spraying with TRIFLUREX may not provide satisfactory control of weeds other than Green Foxtail (Wild Millet) in wheat and barley.

PRECAUTIONS

Warning - skin and eye irritant. Harmful if swallowed or absorbed through the skin. Avoid contact with skin, eyes or clothing. Gloves and goggles should be worn when handling. Avoid inhalation of vapours. Keep out of reach of children. Do not store near heat or flame. Avoid contamination of food stuffs, feed and fish ponds.

FIRST AID

In case of contact with eyes, flush with water immediately. If swallowed do not induce vomiting but contact physician immediately, taking product label with you.

TOXICOLOGICAL INFORMATION: Contains petroleum distillates.

((PAMPHLET))

TRIFLUREX 40 EC

CONTAINS TRIFLURALIN

AGRICULTURAL

HERBICIDE

Directions for Prairie Provinces

A Selective Herbicide for Pre-emergence Weed Control in
Many field Crops, Vegetables and Shelterbelts.

READ THE LABEL AND ATTACHED INSTRUCTIONS BEFORE USING.

GUARANTEE:

Trifluralin.....400 g/L

REGISTRATION NO 21816
PEST CONTROL PRODUCTS ACT

CAUTION POISON

11.3 Litres

Manufactured by: SHAMROCK AGRO-TECH INC:

for:AGROCORP INC.
Brandon, Manitoba

INSTRUCTIONS FOR USE

Carefully read these instructions fully before using TRIFLUREX. Usage not in accordance with these instructions may result in inadequate weed control and may damage crop.

TRIFLUREX is a pre-emergence herbicide which is sprayed on and then mechanically mixed or incorporated into the soil. It kills weeds as they germinate. It does not kill established weeds. TRIFLUREX gives long lasting control of seedlings of annual grasses and broadleaf weeds detailed below.

WARNING

Special attention must be paid to the following aspects of usage

- Rates must be accurately applied
- Apply only on soil types indicated (this excludes peat and muck soils).
- Apply only to soils with less than 15% organic matter which are dry and in good working condition. Soils with the potential of becoming waterlogged must not be treated
- Incorporation is an essential part of application. When correctly applied to crops under normal conditions. TRIFLUREX will not harm the crop. Any stress factors such as wet soils, incorrect planting depth, seedling disease, low temperatures, excessive salt in the soil or drought, could bring about damage to the crop from TRIFLUREX.
- It is recommended that the following crops not follow TRIFLUREX in the rotation cycle: Oats, Small seeded annual grasses such as Canary Seed Grass; Sugarbeets; Creeping Red Fescue; Timothy Grass. Under normal conditions there will not be a carryover However, the above recommendation is precautionary.

DIRECTIONS FOR USE ON OILSEED CROPS

LAND PREPARATION

Cultivate to destroy existing weeds before TRIFLUREX application. When applying to stubble fields, chop and thoroughly mix crop residues into soil for a depth of 10-15 cm. Disc type implements provide the best results. See table below for weeds controlled and crops registered.

APPLICATION

Add the recommended amount of TRIFLUREX 40 EC to clean water in the spray tank during the filling operation and agitate thoroughly. Apply in at least 100 Litres of water per hectare (broadcast basis) using any properly calibrated low-pressure herbicide sprayer that will apply the spray uniformly. Check the sprayer daily to insure proper calibration and uniform application. Apply with ground equipment only. Apply TRIFLUREX to soil surface and incorporate in the same operation if possible. Ensure that large clods are broken. Do not delay incorporation more than 8 hours after application. When applied as directed, TRIFLUREX will not harm crops that follow in rotation; however, over-calibration, non-uniform application or other excess applications may cause injury to crops that follow in rotation.

INCORPORATION

To incorporate, work TRIFLUREX into the soil in two different directions. Use a tandem disc, discer or field (Vibrashank type) cultivator for the first incorporation, set to work 7.5-10 cm deep. Disc or cultivate in cross direction at the same depth. Operate disc implements at 6.5-10 km/hr, cultivators at 10-13 km/hr. TRIFLUREX should not be incorporated with field cultivator when soil is crusted, lumpy or too wet for good mixing action. Disc type implements are preferred on stubble.

Incorporation with implements set to cut less than 7.5 cm deep or more than 10 cm deep may result in erratic weed control or crop damage. Rod weeders, harrows or hoe drills will not properly incorporate TRIFLUREX.

FALL APPLICATION

Check application rates and incorporation instructions above. Ensure timely incorporation. Apply between September 1st and before soil freeze-up for weed control the following year. Two incorporations in the fall are recommended to be followed by shallow tillage (5-8 cm) in the spring before planting. However, if this is not convenient, the first incorporations within 8 hours of spraying can be followed by a second in the spring when the seed bed is being prepared. Always ensure incorporation to the correct depth. (See Spring tillage recommendations below).

SUMMER APPLICATION

Check application rates and incorporation instructions above. Ensure timely incorporation. Apply TRIFLUREX to summer fallow between June 1st and September 1st. Ensure that incorporation must be within 8 hours of spraying. The timing of subsequent incorporations can be suited to the necessity to destroy resistant weed growth later in the season. Before planting in the spring, ensure shallow tillage (5-8 cm). (See Spring tillage recommendations below).

SPRING TILLAGE FOR SEEDBED PREPARATION

In spring, once the soil is warm enough to promote germination, it must be tilled if summer or fall application of TRIFLUREX have been made. Till to a depth of 5-8 cm with a suitable implement such as a disc or Vibrashank cultivator. Follow accepted practices and prevent transplanting weed seedlings.

WEEDS CONTROLLED

The following weeds are controlled with TRIFLUREX when sprayed according to these instructions:

ANNUAL GRASSES

Wild Oats*
Green & Yellow Foxtail (Wild Millet)
Barnyardgrass
Crabgrass
Bromegrass
Cheat
Stinkgrass
Goosegrass
Annual Bluegrass
Persian Darnel

ANNUAL BROADLEAVES

Wild Buckwheat*
Cowcockle*
Pigweed
Lambsquarters
Russian Thistle
Chickweed
Purslane
Knotweed
Carpetweed

NOTE: *Wild Oats and Wild Buckwheat are not fully controlled but retarded by TRIFLUREX. Crop growth will not be seriously hampered by these retarded weeds.

CROPS REGISTERED

Rapeseed	Sunflower
Dry Beans (Field or Kidney Beans)	Mustard
Black Beans	Faba Beans (Horsebeans)
Peas (Field or Canning)	Crambe
Transplanted Shelterbelts	

TRIFLUREX APPLICATION WITH FLUID FERTILIZERS

TRIFLUREX may be mixed with liquid nitrogen fertilizer, (eg. 28-0-0) for application either preplant soil incorporated or postplant incorporated. TRIFLUREX may also be tank mixed with *Avadex BW and liquid fertilizer for postplant incorporated application in wheat or barley. All TRIFLUREX label recommendations regarding rates per hectare, approved crops, incorporation, special instructions, cautions and special precautions must be followed. TRIFLUREX should be poured directly into the liquid fertilizer, mixed thoroughly and applied as soon as possible. Constant agitation is needed until application is complete. Apply the TRIFLUREX/fertilizer mixture with a properly calibrated sprayer. Ensure that the material is applied uniformly to the soil surface. Follow normal TRIFLUREX incorporation procedures.

COMPATIBILITY TEST: Because of the wide range of fertilizer products, small

amounts of liquid fertilizer and TRIFLUREX should be tested for compatibility prior to tank mixing.

- (1) Place 500mL of liquid fertilizer in a 1 L jar
- (2) Pour in 10 mL of TRIFLUREX
- (3) Close jar and shake well
- (4) Leave jar sit for 5 minutes

TRIFLUREX and fertilizer are compatible if TRIFLUREX remains in suspension. If the TRIFLUREX settles out quickly but goes into suspension easily upon shaking the mixture may be used provided constant agitation is utilized.

RECOMMENDED RATES OF TRIFLUREX 40 EC PER HECTARE

CROP	SOIL TEXTURE		REMARKS
	Light	Medium or Heavy	
	Sand Sandy Loam (Less than 6% Organic matter)	Loam Silt Loam Silt Sandy clay loam (All soils with 6-15% organic matter) Silty clay Clay loam Clay	
SPRING APPLICATION	RATE L/ha		
Rapeseed Sunflower Mustard Faba Beans Crambe Peas (Field & Canning) Black Beans Dry Beans (Field & Kidney)	2.0 litres	2.75 litres	TRIFLUREX can be applied immediately prior to, or up to 3 weeks before planting.

Shelterbelts (American Elm, Coragana, Green Ash, Scotch Pine and Siberian Elm	5.5 litres	11.0 litres	Apply prior to transplanting seedlings. The higher rate of application will result in better weed control and longer persistence in the soil.
FALL APPLICATION			
Rapeseed Sunflower Mustard Peas Faba Beans Black Beans Dry Beans (Field & Kidney)	2.75 litres	3.50 litres	Apply and incorporate between September 1st and up to soil freeze-up.
SUMMER APPLICATION			
Rapeseed	All soils at 4.25 litres		Apply and incorporate between June 1st and September 1st. Second and subsequent incorporations may be done any time up to soil freeze-up.

Note: Do not delay incorporation more an 8 hours after application

CONTROL OF GREEN FOXTAIL (WILD MILLET) IN WHEAT & BARLEY

MODE OF ACTION

Most Green Foxtail seeds germinate in the upper 5cm of the soil. TRIFLUREX sprayed and incorporated into this zone will prevent germination. Wheat and Barley seed should be sown below this layer (5-8 cm) to prevent crop damage. Apply and incorporate TRIFLUREX after planting but before emergence of Wheat or Barley.

RECOMMENDED RATES: Light Soils Heavy Soils
 1.4 L/ha 2.1 L/ha

This label transcript service is offered by the Pest Management Regulatory Agency to provide efficient searching for label information. This service and this information do not replace the official hard-copy label. The PMRA does not provide any guarantee or assurance that the information obtained through this service is accurate, current or correct, and is therefore not liable for any loss resulting, directly or indirectly, from reliance upon this service.

+))