

96.10.25

REDUCED 10%

QUAM

DETERGENT/DISINFECTANT

CLEANER DISINFECTANT

DEODORISER SANITIZER

DETERGENT FUNGICIDE

MILDEWSTAT

CAUTION CORROSIVE

GUARANTEE

n-Alkyl (5% C12, 60% C14, 30% C16, 5% C18)
dimethyl benzyl ammonium chlorides.....1.6%
n-Alkyl (5% C12, 60% C14)
dimethyl ethylbenzyl ammonium chlorides.....1.6%

D.I.N. # 01935313

DANGER CORROSIVE TO EYES

COMMERCIAL

ONLY FOR SALE TO, USE AND STORAGE BY SERVICE PERSONS

READ THE LABEL BEFORE USING

REGISTRATION NO. 22192 PEST CONTROL PRODUCTS ACT

NET CONTENTS: L BATCH #

ADVANCE LABORATORIES LTD.
Eastern Shore Ind. Park, Box 130 Halifax Co. N.S.
B0J 1N0
PH: 827-3339
FAX: 827-3773

NOTICE TO USER:

This control product is to be used only in accordance with the directions on this label. It is an offence under the Pest Control Products Act to use a control product under unsafe conditions.

QUAM detergent/disinfection has been designed specifically for hospitals, nursing homes, schools, food processing plants, food service establishments and other institutions where housekeeping is of prime importance.

DISINFECTION- To disinfect hard inanimate surfaces (such as walls, floors, tables), Apply solution with mop cloth, sponge so as to wet all surfaces thoroughly. Allow to remain wet for 10 minutes and then let air dry.

DILUTION RATES- General disinfection- add 15 mL QUAM per L of water. Hospital disinfection 24 mL QUAM per L. of water.

To disinfect toilets, flush toilet, add 75 mL QUAM detergent/disinfectant directly to the bowl water. Swab the bowl completely using a scrub brush or toilet mop, making sure to get under the rim. Let stand 10 minutes and flush.

For heavily soiled areas precleaning is required and recommended. Prepare a fresh solution for each use.

MILDEWSTAT- To control mold and mildew on pre-cleaned, hard non-porous surfaces (such as floors, walls, tables), add 15 mL QUAM per L of water. Apply to surface with cloth, mop, or sponge making sure to wet all surfaces completely. Let air dry. Prepare a fresh solution for each use. Repeat application at weekly intervals or when mildew growth reappears.

SANITIZATION- To sanitize previously cleaned and rinsed non-porous food contact surfaces, prepare 200 ppm active quaternary QUAM solution by adding 6 mL QUAM per L of water. To sanitize immovable items (such as tanks, chopping blocks, counter tops), flood the area with 200 ppm active quaternary for at least 60 seconds making sure to wet all surfaces completely for at least 60 seconds. Let air dry. Prepare a fresh solution for each use. To sanitize mobile items (such as drinking glasses, eating utensils) immerse in 200 ppm QUAM solution for at least 60 seconds making sure to immerse completely. Remove and let air dry. Prepare a fresh solution daily or more frequently as soil is apparent.

PRECAUTIONARY STATEMENTS: DANGER HAZARD TO HUMANS AND DOMESTIC ANIMALS Keep out of reach of children. Causes severe eye and skin damage. DO not get in eyes or skin or clothing. Wear goggles or face shield and rubber gloves when handling. Harmful

or fatal if swallowed. Avoid contamination of food.

FIRST AID- In case of contact immediately flush eyes or skin with plenty of water or at least 15 minutes. for eyes call a physician. remove and wash contaminated clothing before reuse. If swallowed drink amounts of water. Avoid alcohol. Call a physician or poison control centre immediately.

TOXICOLOGICAL INFORMATION- Probable mucosal damage may contraindicate the use of gastric lavage. Measurable against circulatory shock, respiratory depression and convulsion may be needed.

QUAM detergent/disinfectant has been designed specifically for hospitals, nursing homes, schools, food processing plants, food service establishments and other institutions where housekeeping is of prime importance in controlling the hazard of cross contamination.

QUAM when diluted at the rate of 24 mL per L is an effective disinfectant for hospital use, when used as directed. QUAM is formulated to disinfect inanimate hard surfaces such as walls, floors, sink tops, toilet bowls, tables, chairs, telephone and bed frames. For larger areas such as operating rooms, patient care facilities and restrooms, the product is designed to provide both general cleaning and disinfections.

In addition QUAM deodorizes those areas which generally are hard to keep fresh smelling, such as garbage storage areas, empty garbage bins and cans, toilet bowls and any other areas which are prone to odours caused by microorganisms.

DISINFECTION- At 15 mL dilution, QUAM exhibits effective disinfectant activity against the organisms: Salmonella choleraesuis, and Staphylococcus aureus. At this level the product is also fungicidal against the pathogenic fungi, Trichophyton mentagrophytes.

MOLD AND MILDEWSTAT- QUAM prevents and controls mold and mildew and the odours they cause on hard non-porous inanimate surfaces at 15 mL per L dilution.

SANITIZATION- When used for sanitization of previously cleaned food equipment or food contact areas active quaternaries of 200 ppm. QUAM is an effective sanitizer when diluted in water up to 500 ppm hardness (CaCO₃).

DISPOSAL:

1. Rinse the emptied container thoroughly and add the rinsings

- to the treatment site.
2. Follow provincial instructions for any required additional cleaning of the container prior to its disposal.
 3. Make the empty container unsuitable for further use.
 4. Dispose of the container in accordance with provincial requirements.
 5. For information on the disposal of unused, unwanted product and the cleanup of spills, contact the Provincial Regulatory Agency or the Manufacturer.

This label transcript service is offered by the Pest Management Regulatory Agency to provide efficient searching for label information. This service and this information do not replace the official hard-copy label. The PMRA does not provide any guarantee or assurance that the information obtained through this service is accurate, current or correct, and is therefore not liable for any loss resulting, directly or indirectly, from reliance upon this service.

+))