

99.03.25

SEVIN (R)

WETTABLE POWDER

INSECTICIDE

10 kg

COMMERCIAL

FOR MIXING AS A SPRAY TO CONTROL EXTERNAL
PARASITES OF POULTRY AND LIVESTOCK

GUARANTEE:

Carbaryl 50% (W/W)

REGISTRATION NO. 9001
PEST CONTROL PRODUCTS ACT

READ THE LABEL BEFORE USING

WARNING

POISON

Sevin®: Registered trademark Rhone Poulenc, Nederland, B.V.

Distributed by
Solvay Animal Health, Inc.,
Kitchener, Ontario N2C 1L4

POULTRY INSECT CONTROL

On chickens, ducks, geese, turkeys and domestic game birds: For northern fowl mite and lice and as a supplement to premise treatment for chicken mite, fleas and fowl ticks: Use 1 kg per 100 litres of water (10 g per litre). Spray at the rate of 5 litres per 100 birds. Direct spraying for chicken mites and fleas is a supplement to spraying roosts and buildings for control of these pests. Repeat in 4 to 8 weeks or as necessary.

In or around poultry houses and farm buildings: For poultry mites, fleas and bed bugs, use 1 kg per 100 litres of water. For fowl ticks, use 4 kg per 100 litres of water. Spray 5 to 10 litres per 100 square meter of wall, bedding, litter or root surfaces. Force spray into cracks. Repeat as needed. Ventilate while spraying. Do not apply to livestock, poultry or poultry nest litter within 7 days of slaughter. Avoid contamination of nests, eggs, feed and water troughs. Do not treat birds with respiratory or other disease that may be irritated by dust.

LIVESTOCK INSECT CONTROL

On beef cattle, horses, sheep, goats and hogs: For lice, hornflies, winter ticks and fleas, use 10 kg per 1000 litres of water and apply thoroughly to entire animal at the rate of 5 litres per head. Repeat as necessary, but not more often than twice weekly. For lice, repeat at 14 to 18 day intervals.

On dairy cattle: For lice, hornflies, winter ticks and fleas, mix as directed above and spray carefully along topline, sides and legs. Avoid direct application to underline and udder. Do not apply before milking: only immediately after milking. Wash udder thoroughly before milking. Do not contaminate milking utensils. Repeat if needed, but not more often than twice weekly. In order to prevent residues in milk or meat, do not apply within 7 days of slaughter and do not exceed dosages given above.

PRECAUTIONS: KEEP OUT OF REACH OF CHILDREN. Harmful if swallowed or inhaled. Avoid contact with skin. Wash thoroughly after handling. Change contaminated clothing daily. Avoid contamination of feed, foodstuffs, feed troughs and water receptacles. Do not contaminate streams, ponds, water supplies. Highly toxic to bees - do not use where bees are foraging.

FIRST AID: If on skin, remove contaminated clothing and wash immediately with soap and water. If in eyes, flush with water for at least 15 minutes. If swallowed, induce vomiting by touching back of throat with finger or blunt object. Get medical attention immediately.

TOXICOLOGY: Carbaryl is a moderate reversible cholinesterase inhibitor. Early symptoms: headache, nausea, vomiting and blurring of vision. This may progress to dizziness, sweating, profound weakness, salivation, chest pain, abdominal cramps and diarrhea. Large doses of atropine may be needed to control symptoms. Full atropinization may be necessary. If no atropine has been given by mouth, intravenous doses of 1 to 2 mg may be given at 15 minute intervals until symptoms are relieved or full atropinization is achieved. If atropine has already been given, additional doses of 1 mg may be given until atropinization is achieved. 2-P.A.M. is contraindicated.

NOTICE TO USER: This control product is to be used only in accordance with the directions on this label. It is an offense under the Pest Control Products Act to use a control product under unsafe conditions.

LOT NO: 3347

This label transcript service is offered by the Pest Management Regulatory Agency to provide efficient searching for label information. This service and this information do not replace the official hard-copy label. The PMRA does not provide any guarantee or assurance that the information obtained through this service is accurate, current or correct, and is therefore not liable for any loss resulting, directly or indirectly, from reliance upon this service.

+))