

85.12.31

GUARDSMAN(R)

CAPTAN 5%

DUST FUNGICIDE

COMMERCIAL

READ THE LABEL BEFORE USING

REGISTRATION NO. 13114 PEST CONTROL PRODUCTS ACT

GUARANTEE: CAPTAN 5%

NET CONTENTS: 20 KG

DISTRIBUTED BY

VAN WATERS & ROGERS LTD.

EDMONTON CALGARY TORONTO VANCOUVER MONTREAL WINNIPEG

P.O. BOX 2009 V6B 3R2

Exposure to Captan may produce long term health effects. To minimize exposure, follow directions outlined on this label. Directions for use: To obtain satisfactory results, use good dusting equipment and apply thoroughly a uniform deposit of dust. May be applied up to 2 days of harvest. Use the indicated amount per hectare.

CUCUMBERS: Anthracnose and Scab - use 34 to 50 kg. on young plants and 50 to 67 kg. on mature plants at 5-7 day intervals. Treat at the first sign of disease.

TOMATOES: Anthracnose, Early and Late Blight, Septoria & GrayLeaf Spot - 34 to 55 kg. at the first sign of infection at 7-10 day intervals. For severe infections or on mature plants, use 70 to 83 kg.

STRAWBERRY: Grey Mold Rot, Leaf Spot - use 50 to 83 kg. Begin application when new growth starts in spring. Repeat at weekly

intervals. Under conditions favourable to fruit rot, continue applications throughout the harvest period. Apply not later than 2 days before each picking.

POTATO SEED PIECE TREATMENT: Brown Eye Disease (Damping-off), Scab, Seed Rot, Rhizoctonia, coat each piece evenly with a light film of dust. Use at the rate of 415 g per 25 kg. Treat within 6 hours of cutting. If planting is to be delayed for a day or two, store in open crates and provide sufficient ventilation to allow cut surfaces to dry. Label all unplanted treated seed: "Poisonous to man and animals. This seed has been treated with captan." Do not sell unplanted seed potatoes for use for food or feed.

KEEP OUT OF REACH OF CHILDREN. Never handle material with bare hands. Use rubber gloves; do not use leather or cloth gloves. Avoid spilling on skin or clothing. Do not breathe spray mist (dust). Wear a respirator and goggles while mixing, loading and spraying. When handling or applying, wear goggles, clothing that completely covers arms and legs to minimize exposure. Change contaminated clothing daily and wash thoroughly before re-use. Wash hands, face and arms thoroughly after handling captan and before drinking, eating or smoking. Shower after use or at the end of the work day. If an accident occurs, remove contaminated clothing and wash skin thoroughly with soap and water. Wash clothing and gloves before re-use. Always work out of the wind during mixing and loading operations. Keep unprotected persons out of operating area or vicinity where there may be drift. Thoroughly wash treated fruits and vegetables prior to eating.

SYMPTOMS OF POISONING FIRST AID: In case of poisoning, call a physician immediately. IF ON SKIN, remove contaminated clothing and wash skin thoroughly with soap and water. IF IN EYES, flush with water for 5-10 minutes and obtain medical attention. IF SWALLOWED, give the patient one to two glasses of water, and cause vomiting by giving one dose of syrup of ipecac. If the patient does not vomit within 20 minutes, give a second dose. If syrup of ipecac is not available, give the patient one to two glasses of water and cause vomiting by inserting finger down the throat. Repeat with water until vomit fluid is clear. The patient should be lying down with the head below the level of the feet. DO NOT TRY TO CAUSE VOMITING IF THE PATIENT IS UNCONSCIOUS OR IN A CONVULSIVE STATE.

NOTICE TO USER: This control product is to be used only in accordance with the directions on the label. It is an offence

under the PEST CONTROL PRODUCTS ACT to use a control product under unsafe conditions.

NOTICE TO BUYER: Seller's guarantee shall be limited to the terms set out on the label and subject thereto the buyer assumes any risk to persons or property arising from the use or handling of this product and accepts the product on that condition.

This label transcript service is offered by the Pest Management Regulatory Agency to provide efficient searching for label information. This service and this information do not replace the official hard-copy label. The PMRA does not provide any guarantee or assurance that the information obtained through this service is accurate, current or correct, and is therefore not liable for any loss resulting, directly or indirectly, from reliance upon this service.

+))