2000/2001

ASSASSIN

KILLS RATS AND MICE

TUE RATS ET SOURIS

ALL-WEATHER/PAR TOUS LES TEMPS

BAIT BLOX/APPAT BLOX

DOMESTIC/DOMESTIQUE

REG. NO./NO. D'ENR. 23670

P.C.P. ACT/LOI P.A.

GUARANTEE/GARANTIE:

DIPHACINONE 0.005%

READ THE LABEL BEFORE USING

(SEE LEFT PANEL FOR ADDITIONAL PRECAUTIONARY ELEMENTS.)

KEEP OUT OF REACH OF CHILDREN

LIRE L'ETIQUETTE AVANT USAGE

(VOIR A DROITE POUR PRECAUTIONS SUPPLEMENTAIRES.)

GARDER HORS DE LA PORTEE DES ENFANTS

NET CONTENTS: 0.8 kg (1.75 lbs.)

CONTENU NET: 0,8 kg (1,75 lbs.)

DISTRIBUTED BY/DISTRIBUE PAR: CITADEL ANIMAL HEALTH/SANTE ANIMALE CAMBRIDGE, ON N3C 2W4

DIRECTIONS FOR USE: Use bait to control the Norway rat,(Rattus norvegicus), Roof rat(Rattus rattus) and House mouse(Mus musculus) in and around dwellings.

RATS: Place 3 to 16 pieces of ASSASSIN BLOX(usually at intervals of 4.5 to 9 meters) per placement. Maintain an uninterrupted supply of fresh bait for 10 days or until signs of rat activity

cease.

HOUSE MICE: Place 1 piece of ASSASSIN BLOX at intervals of 2.5 to 3.5 meters per placement. Two pieces may be needed at points of high mouse activity. Maintain an uninterrupted supply of fresh bait for 15 days or until signs of mouse activity cease. Replace contaminated or spoiled bait immediately. Collect and dispose of unconsumed bait and dead exposed animals according to local requirements. To prevent reinfestation, eliminate food, water, and harborage as much as possible. If reinfestation does occur, repeat treatment. Where a continuous source of infestation is present, establish permanent bait stations and replenish bait as needed.

PRECAUTIONS: Keep away from humans, domestic animals and pets. Avoid contact with eyes and clothing. Wash thoroughly with soap and water after using. If swallowed, this material may reduce the clotting ability of the blood and cause bleeding.

FIRST AID: If in eyes, flush with water for 15 minutes. If on skin, wash with soap and water. If swallowed, give patient one or two glasses of water and induce vomiting by touching back of throat with a finger or a blunt object; repeat until vomit fluid is clear. Do not give anything by mouth to an unconscious person. Contact a physician or Poison Control Centre immediately.

TOXICOLOGICAL INFORMATION: If ingested, Vitamin K₁ in the form of intravenous, intramuscular or subcutaneous injections, or by oral ingestion are suggested remedial treatments for anticoagulant poisoning. The severity of the case measured by establishing prolonged prothrombin times(P.T.) will determine appropriate therapy. Monitoring P.T. will indicate the necessity of repeated treatments.

ENVIRONMENTAL HAZARD: Keep out of lakes, streams or ponds.

USE LIMITATIONS: Treated baits must be placed in locations not accessible to children, pets, domestic animals and wildlife, or in tamperproof bait stations. Do not place bait in areas where there is a possibility of contaminating food or surfaces that come in direct contact with food.

SELECTION OF TREATMENT AREAS: Determine areas where rats and/or mice will most likely find and consume the bait. Generally, these areas are along walls, by gnawed openings, in or beside burrows, in corners and concealed places, between floors and walls, or in locations where rodents or their signs have been observed. Remove as much food as possible.

DISPOSAL: Do not reuse empty container. Wrap and dispose of empty container in household garbage. Dispose of unconsumed bait according to Local requirements.

RODENT DISPOSAL: Dispose fo dead rodents in garbage or by burying.

This label transcript service is offered by the Pest Management Regulatory Agency to provide efficient searching for label information. This service and this information do not replace the official hard-copy label. The PMRA does not provide any guarantee or assurance that the information obtained through this service is accurate, current or correct, and is therefore not liable for any loss resulting, directly or indirectly, from reliance upon this service.

+))