

85.12.31

TOTIL 19-500

(PRODUCT OF THE UNITED STATES)

COMMERCIAL BENZALKONIUM CHLORIDE SOLUTION

(CALGON STANDARD)

STAPHYLOCIDAL/PSEUDOMONACIDAL/SALMONELLACIDAL/DISINFECTS

DEODORIZES/SANITIZES

CAUTION

POISON

DANGER

CORROSIVE

READ THE LABEL BEFORE USING

REGISTRATION NO.: 14898

PEST CONTROL PRODUCTS ACT

GUARANTEE

n-alkyl(40% C12, 50% C14, 10% C16) dimethyl benzyl ammonium chloride.....	10.0%
Tetrasodium ethylenediamine tetraacetate.....	0.8%
Ethyl Alcohol.....	2.5%

DIN 429848

MANUFACTURED BY - CALGON CORPORATION  
COMMERCIAL DIVISION  
7501 PAGE AVENUE - ST. LOUIS, MISSOURI  
63166  
FOR - CALGON CANADA, INC.  
27 FINLEY ROAD  
BRAMALEA, ONTARIO  
L6T 1B2

CONCENTRATE MUST BE DILUTED BEFORE USE  
FOR MANUAL APPLICATION ONLY

DIRECTIONS FOR USE:

LIGHTLY DUTY CLEANING AND SANITIZATION: 2 mL of Totil added to 1 liter of water will provide 200 ppm active quaternary germicide. Apply solution and allow to soak on surface for 10 minutes before rinsing. Rinse food preparation surfaces with potable water before reuse.

HEAVY DUTY CLEANING AND DISINFECTION: 8 mL of Totil added to 1 liter of water will provide 780 ppm active quaternary germicide. Follow instructions for Light Duty Cleaning and Sanitization. For disinfection of surfaces heavily covered with dirt or organic material first clean then apply at the rate for disinfection.

PRECAUTIONS:

DANGER: KEEP OUT OF REACH OF CHILDREN. Corrosive to the eye. Do not use Totil with soap or other detergents. Rinse empty container thoroughly with water and discard it. Do not get in eyes, on skin or on clothing. Protect eyes and skin when handling. Harmful if swallowed. Avoid contamination of food.

FIRST AID: In case of contact, immediately flush eyes or skin with plenty of water for at least 15 minutes. For eyes, call a physician. Remove and wash contaminated clothing before reuse. If swallowed, drink promptly a large quantity of milk, egg whites, gelatin solution or, if these are not available, drink large quantities of water. Call a physician immediately.

NOTE TO PHYSICIAN: Probable mucosal damage may contraindicate the use of gastric lavage. Measures against circulatory shock, respiratory depression and convulsion may be needed.

NOTICE TO USER:

This control product is to be used only in accordance with the directions on this label. It is an offence under the Pest Control Products Act to use a control product under unsafe conditions.

\*\*\*\*\*

This label transcript service is offered by the Pest Management Regulatory Agency to provide efficient searching for label information. This service and this information do not replace

the official hard-copy label. The PMRA does not provide any guarantee or assurance that the information obtained through this service is accurate, current or correct, and is therefore not liable for any loss resulting, directly or indirectly, from reliance upon this service.

+) )