

Nufarm

AMIZOL SOLUBLE POWDER FOR WEED CONTROL

READ THE LABEL BEFORE USING

INDUSTRIAL USE

FOR WEED CONTROL IN INDUSTRIAL AND OTHER NON-CROPPED AREAS

GUARANTEE: Amitrol.....90%

REGISTRATION NO. 7559 PEST CONTROL PRODUCTS ACT

CAUTION

POISON

NET CONTENTS: 22.7 KG

Nufarm Agriculture Inc.  
5507 1<sup>st</sup> Street SE  
Calgary, Alberta T2H 1H9

**DIRECTIONS FOR USE: AMIZOL FOR WEEDS**

Apply Amizol anytime weeds are green and actively growing. Weeds treated after they have developed strong stems may remain standing for the rest of the season, even though they are dead. These standing weeds are unsightly and a fire hazard. Therefore, for most effective control of tall weeds, cut and remove the weeds if possible, then spray the regrowth. For best results, treat when soil moisture is adequate for rapid weed growth. Use enough water to thoroughly wet weed foliage--500 to 1000 litres per hectare. In highway spraying, a strip 2.5 metres wide and 2.0 kilometres long is about 1/2 hectare. Be sure Amizol is completely dissolved before spraying. Do not use for selective weed control in turf or edible crops.

**Some of the Weeds Controlled Include:**

Alfalfa, Blue grasses, Canada thistle, Carpetweed, Catch fly, Chickweed (common), Chrysanthemum weed, Cocklebur, Crab grass, Cough grass, Dandelion, Downy brome grass, Foxtail (wild millet), Goose grass, Groundsel, Hemp nettle, Jimson weed, Knotweed, Kochia, Lambsquarters, Mustard, Nightshade, Pigweed, Plantain, Poison Ivy, Purslane, Ragweed, Shepherd's purse, Smartweeds, Sow thistle, Stinkweed, Velvet grass,

White cockle, Wild buckwheat, Wild oats, Witchgrass, Yellow rocket.

**Amounts per Hectare**

A. Control or suppress annual grasses and broadleaf weeds where residual control is not required. Spray when most weeds are 8 to 10 cm tall.

AMIZOL: 2.25 kg      Water: 1000 L

B. Control annual grasses, annual and perennial broadleaf weeds, suppress perennial grasses. Spray when most weeds are 15-25 cm tall.

AMIZOL: 4.5 kg      Water: 1000 L

C. Cattails: treat after catkins are fully formed until frost.  
Ground application

AMIZOL: 10-11 kg      Water: 3000 L

D. Poison ivy, poison oak. Treat after leaves are fully developed in the spring until leaves start to turn colour in the fall.

AMIZOL: 2.25 kg      Water: 1000 L

**DIRECTIONS FOR USE: AMIZOL FOR WOODY PLANTS**

Use Amizol in a right-of-way maintenance program as a special tool for cleaning up spot infestations of the species listed below. Most of these are killed to ground line by foliage sprays of 2,4-D, but resprout badly the next year. Amizol is a good translocator and kills most of the roots as well as the tops. Amizol is specific on the species listed. It should not be expected to kill other species. Use the standard foliage stem method for high volume sprays along fencerows, highways, and utility rights-of-way. Spray all foliage and stems of plants from the time foliage is fully developed until plants begin to go dormant. For effective kills, all leaves, stems, and suckers must be thoroughly wet to ground line.

**Kg of Amizol to Use per  
1000 litres of water**

SPECIES	Salmonberry	4	Add 250 mL of spreader sticker per 1000 L water.
	Western dewberry (blackberry)	4	
	Bigleaf maple	4	
	Honeysuckle	4	
	Locust	3 to 4	
	Ash	4 to 6	
	Sumac	3 to 4	

**DO NOT APPLY BY AIR**

**PRECAUTIONS:** KEEP OUT OF REACH OF CHILDREN. Avoid spray-drift as this product may injure beans, peas, tomatoes, some ornamentals, etc. (Coarse sprays are less likely to drift). DO not use same sprayer for other purposes unless thoroughly cleaned. To clean sprayer, rinse all parts of sprayer with water several times. Do not store near fertilizers, seeds, insecticides, or fungicides or use in a greenhouse. Do not contaminate water used for irrigation or domestic purposes. Avoid contact with skin, eyes or clothing.

**FIRST AID:** Remove skin contamination by washing with soap and water. **For eyes,** flush immediately with clear water. Do not take internally. **If swallowed,** INDUCE VOMITING by touching the back of the throat with finger or blunt object. Get medical attention or contact a poison control centre IMMEDIATELY. Take container, label or product name and Pest Control Product Registration Number with you, when seeking medical attention.

**TOXICOLOGICAL INFORMATION:** This product may affect thyroid function. Overexposure can be determined by monitoring thyroid function. If applicator is under treatment for hypo- or hyperthyroidism, they should be removed from exposure until the euthyroid state is achieved.

**DISPOSAL:**

1. Triple- or pressure-rinse the empty container. Add the rinsings to the spray mixture in the tank.
2. Follow provincial instruction for any required additional cleaning of the container prior to its disposal.
3. Make the empty container unsuitable for further use.
4. Dispose of the container in accordance with provincial requirements.
5. For information on disposal of unused, unwanted product, contact the manufacturer or the provincial regulatory agency. Contact the manufacturer and the provincial regulatory agency in case of a spill, and for clean-up of spills.

**NOTICE TO USER:** This control product is to be used only in accordance with the directions on this label. It is an offence under the PEST CONTROL PRODUCTS ACT to use a control product under unsafe conditions.

**NOTICE TO BUYER:** Seller's guarantee shall be limited to the terms set out on the label and, subject thereto, the buyer assumes the risk to persons or property arising from the use or handling of this product and accepts the product on that condition.