

88.10.28

CLEANER - DISINFECTANT

FOR HOG, CATTLE AND POULTRY BUILDINGS

F O R M U L A 9

NETTOYEUR - DISINFECTANT

POUR BATIMENTS DESTINES AUX BESTIAUX ET VOLAILLE

FOR AGRICULTURAL USE ONLY

POUR USAGE AGRICOLE UNIQUEMENT

GUARANTEE:

N-Alkyl (5% C12, 60% C14, 30% C16, 5% C18)
dimethyl benzyl ammonium chloride..... 0.8%
N-Alkyl (68% C12, 32% C14) dimethyl
ethyl benzyl ammonium chloride..... 0.8%

GARANTIE:

Chlorure N-Alkyl (5% C12, 60% C14, 30% C16,
5% C18) dimethyl benzyl ammonium..... 0.8%
Chlorure N-Alkyl (68% C12, 32% C14)
dimethyl ethyl benzyl ammonium..... 0.8%

CAUTION - ATTENTION

CORROSIVE - CORROSIF

COMMERCIAL

READ THE LABEL BEFORE USING

LIRE L'ETIQUETTE AVANT L'EMPLOI

REGISTRATION NO. 20245

PEST CONTROL PRODUCTS ACT

NO. D'ENREGISTREMENT 20245

LOI SUR LES PRODUITS ANTIPARASITAIRES

Contenu Net / Net Contents Litres

Lot #

DELL TECH LABORATORIES LTD.
13 ROUTLEDGE STREET,
HYDE PARK, ONTARIO
NOM 1Z0

DESCRIPTION: Formula 9 is a heavy-duty cleaner with disinfectant properties. Good removal of soil load is an important part of all surface disinfection and Formula 9 contains detergents and builders to give excellent cleaning. Unlike products designed for disinfection at lower concentration use (eg. 250 to 1) Formula 9 delivers both cleaning and disinfectant punch at recommended use dilution. Formula 9 will penetrate through organic soil and allow complete cleaning and disinfection.

DIRECTIONS FOR USE: General Cleaning and Disinfection of Livestock and Poultry Buildings: Remove all livestock and poultry from buildings. Remove heavy accumulations of manure and soil from floors, walls, pens, equipment using shovel and brush. Apply "Formula 9" at 1 part per 25 parts hot or cold water. Allow 10-15 minutes for solution to penetrate and loosen manure and organic soil. Rinse thoroughly using high pressure hot or cold water. Repeat cleaning and rinsing steps if necessary.

Sanitizing: To sanitize pre-cleaned, rinsed, non-porous surfaces, prepare a Sanitizing solution of 1 part "Formula 9" to 80 parts water and saturate all surfaces. Allow to air dry, before housing poultry or livestock.

Equipment Cleaning and Disinfection: Empty all troughs, racks, etc. Apply "Formula 9" at 1 part per 25 parts hot or cold water. Allow 10-15 minutes contact time during which scrubbing or brushing can be applied. Forks, shovels, scrapers and handling equipment should be immersed in the cleaning solution. Thoroughly pressure rinse all surfaces using hot or cold water. All direct food contact surfaces such as feeders and waterers must be rinsed with potable water prior to use.

Note: At 1 part to 25 parts water "Formula 9" exhibits effective disinfectant activity against: *Escherichia coli*, *Salmonella choleraesuis*, *Salmonella pullorum*, *Staphylococcus aureus*. At this level the product is also fungicidal against *Aspergillus niger*. At 1 part to 21 parts water "Formula 9" demonstrates activity against *Pseudomonas aeruginosa* and *Pseudomonas fluorescens*. Efficacy tests have demonstrated that this product

is an effective bactericide, fungicide and virucide in the presence of organic soil (5% blood serum). "Formula 9" is an effective sanitizer when diluted with water up to 250 ppm hardness (CaCO₃).

PRECAUTION: Keep out of reach of children. Causes eye and skin damage. Do not get in eyes, on skin or clothing. Harmful if swallowed. Wear glasses or face shield and rubber gloves when handling. Avoid contamination of food. When disinfecting surfaces coming in direct contact with food, rinse thoroughly with potable water.

FIRST AID INSTRUCTIONS: In case of contact, immediately flush eyes or skin with plenty of water for at least 15 minutes. For eyes, contact a physician. Remove and wash all contaminated clothing before re-use. If swallowed, drink milk, egg whites, gelatin solution or if none of these are available, drink large quantities of water. Call a physician immediately.

NOTICE TO USER: This control product is to be used only in accordance with the directions on this label. It is an offense under the Pest Control Products Act to use a control product under unsafe conditions.

TOXICOLOGICAL INFORMATION: Probable mucosal damage may contraindicate the use of gastric lavage. Measures against circulatory shock, respiratory depression and convulsion may be needed.

DISPOSAL:

1. Rinse the emptied container thoroughly and add the rinsings to treatment site.
2. Follow provincial instructions for any required additional cleaning of the container prior to its disposal or reconditioning.
3. Dispose of the container in accordance with provincial requirements.
4. For information on the disposal of unused, unwanted product and the cleanup of spills, contact the regional office of Conservation and Protection, Environment Canada.
5. For small spills. Recover free liquid. Soak up residue with solids absorbent. DO NOT FLUSH INTO SURFACE STREAMS. INFORM ENVIRONMENT CANADA AS IN #4.

This label transcript service is offered by the Pest Management Regulatory Agency to provide efficient searching for label

information. This service and this information do not replace the official hard-copy label. The PMRA does not provide any guarantee or assurance that the information obtained through this service is accurate, current or correct, and is therefore not liable for any loss resulting, directly or indirectly, from reliance upon this service.

+))