

85.12.31

Read the Label Before Using

STANCHEM

MCPA
SODIUM 48 SOLUTION

WEEDKILLER FOR CEREALS

AGRICULTURAL

WARNING POISON

GUARANTEE:

MCPA (present as a Sodium Salt)
300 grams per litre

REGISTRATION NO.7812
PEST CONTROL PRODUCT ACT

A DIVISION OF PPG INDUSTRIES CANADA LTD.
60 TITAN ROAD
TORONTO, ONTARIO
M8Z 2J8

MONTREAL WINNIPEG EDMONTON CALGARY

PRECAUTIONS

1. This product may damage susceptible crops. Do not use near susceptible crops such as rapeseed, mustard, tomatoes, grapes, many vegetables and ornamentals.
2. Do not apply on a windy day.
3. Avoid creating spray drift. (Use a coarse spray, low pressure and higher volume of water to reduce this problem).

4. Do not use spray equipment used for applying this product for spraying susceptible crops.
5. Avoid contamination of water supplies, especially those used for irrigation.
6. Avoid contact with skin, eyes or clothing.

FIRST AID

In case of skin contact, wash immediately with soap and water and remove contaminated clothing. If skin irritation occurs obtain medical attention. In case of eye contact, flush with plenty of water and obtain medical attention. If swallowed induce vomiting. If patient is unconscious give him air; call a physician or poison control centre.

CONTAINER DISPOSAL

Do not reuse container. Rinse out container with water after use, destroy and dump. Do not contaminate water supplies with rinse water.

STORAGE

Store away from fertilizers, seed, insecticides, fungicides and other herbicides.

MCPA SODIUM 48 SOLUTION

MIXING INSTRUCTIONS

Calculate the number of hectares the sprayer will treat with one filling. Determine the amount of STANCHEM MCPA SODIUM 48 SOLUTION to be used in the sprayer by multiplying the number of hectares to be treated by the recommended application rate per hectare. Add one-half the required amount of water to the sprayer tank. Add the correct amount of STANCHEM MCPA SODIUM 48 SOLUTION. Mix well, then add the remainder of the water. Continue mixing until spray mixture is uniform.

NOTICE TO BUYER

Seller's guarantee shall be limited to the terms set out on the label and subject thereto. The buyer assumes the risk to persons or property arising from the use or handling of this product and accepts the product on that condition.

NOTICE TO USER

THIS CONTROL PRODUCT IS TO BE USED ONLY IN ACCORDANCE WITH THE DIRECTIONS ON THIS LABEL. IT IS AN OFFENCE UNDER THE PEST CONTROL PRODUCTS ACT TO USE A CONTROL PRODUCT UNDER UNSAFE CONDITIONS.

MCPA SODIUM 48 SOLUTION
Weedkiller for Cereals

STANCHEM MCPA SODIUM 48 SOLUTION control many broad-leaved weeds in cereals and number of other crops. Before using - select the Application Rate, follow the Mixing Directions (See other panel) and observe the Precautions (See other panel).

DIRECTIONS FOR USE

CEREALS - Apply a minimum of 40 L/ha.

CROP	TIMING	WEEDS AND RATES PER HECTARE	
		SUSCEPTIBLE WEEDS	MORE RESISTANT WEEDS
Spring Wheat, Barley, Oats (see Note 3)	2 leaf stage to early shot-blade stage. (See Note 1)	1.5-2.0 L	2.0-3.5 L
Winter Wheat, Fall Rye	Early Spring to shot-blade stage.	1.5-2.0 L	2.0-3.5 L (see Note 2)

COMMON SUSCEPTIBLE WEEDS - Flixweed, Kochia, Lamb's Quarters, Mustard, Ragweed, Shepherd's Purse, Stinkweed.

COMMON MORE RESISTANT WEEDS - Canada Thistle (top kill only), Dog Mustard, Pigweed, Russian Thistle, Sow Thistle (top kill only).

- NOTE 1. Spray at the 4 leaf stage for best crop tolerance and weed control.
- NOTE 2. Rates used to control more resistant weeds may cause some crop injury. Improved yields from better weed control will usually more than offset this injury.
- NOTE 3. For weed control in cereals underseeded with certain legumes, follow the recommendations in the Forage Crop and Underseeded Cereals Section.

FLAX

The crop must be a minimum of 5 cm high. Application must be made before the crop reaches the early bud stage. Use 100-150 litres water per hectare to reduce risk of damage. Apply 1.5-2.0 litres per hectare to control susceptible weeds. If more resistant weeds are abundant, apply 2.0-3.5 litres per hectare. Rates of application of 1.75 litres per hectare or more may cause some delay in maturity and some crop injury. The crop injury will likely be more than offset by increased yields due to effective weed control.

FORAGE CROPS AND CEREALS UNDERSEEDED WITH LEGUMES

Do not use on Sweet Clover.

For underseeded cereals, follow the recommendations for legumes below.

CROP	TIMING	MCPA SODIUM 48 SOLUTION PER HECTARE	WATER PER HECTARE
Seedling Alfalfa, Red and Alsike Clover and White Dutch Clover.	Apply from first to third trifoliolate leaf stage. Do not spray after the crop is 10 cm high.	1.2-1.5 L (see Note)	170-220 L

NOTE: Some injury may occur to legumes. This will be reduced by using the recommended volume of water.

GRASS CROPS AND CEREALS UNDERSEEDED WITH GRASSES

For underseeded cereals, follow the recommendations volume of water.

CROP	TIMING	MCPA SODIUM 48 SOLUTION PER HECTARE	WATER PER HECTARE
Seedling Grasses	Apply from 3 leaf stage to just before shot-blade stage.	1.75 L	45 or more litres

Established Grasses	Before shot-blade stage in Spring or after harvest. Do not graze treating pasture within 7 days of spraying.	1.75-3.75 L	170 L
---------------------	--	-------------	-------

PEAS

Treat the crop when the vine length is 10 to 20 cm. Use 110 to 220 litres per hectare and the higher amount is preferable. Apply 0.5 to 1.5 litres per hectare MCPA SODIUM 48 SOLUTION to control susceptible weeds.

This label transcript service is offered by the Pest Management Regulatory Agency to provide efficient searching for label information. This service and this information do not replace the official hard-copy label. The PMRA does not provide any guarantee or assurance that the information obtained through this service is accurate, current or correct, and is therefore not liable for any loss resulting, directly or indirectly, from reliance upon this service.

+))