

Notification Change

Bayer Cropscience Inc. □



ROVRAL FLO

FUNGICIDE

CONTAINS IPRODIONE

FOR CONTROL OF DISEASES IN CANOLA

AGRICULTURAL

GUARANTEE: IPRODIONE.....250g/L

REGISTRATION NO. 18977
PEST CONTROL PRODUCTS ACT

READ THE LABEL AND BOOKLET BEFORE USING.



CAUTION

POISON

WARNING: SKIN AND EYE IRRITANT

Notification Change

~~Aventis CropScience Canada Co.
295 Henderson Drive
Regina, Saskatchewan
S4N 6C2~~

~~Telephone: 1-888-AVENTIS (1-888-283-6847)~~

**Bayer Cropscience Inc. □
Suite 100 □
3131-114th Ave. S.E. □
Calgary, AB T2Z 3X2**

NET Contents: 8, 10 LITRES

IN CASE OF SPILLS, POISONING, OR FIRE TELEPHONE EMERGENCY RESPONSE
NUMBER, COLLECT 1-800-334-7577 (24 HOURS A DAY)

~~Aventis~~ and ROVRAL® are registered trademarks of the ~~Aventis Group~~.

USE RESTRICTIONS:

Do not store ROVRAL FLO near foodstuffs.
Thoroughly clean application equipment after use.
Do not contaminate domestic or irrigation waters.
Do not allow spray drift onto oats.

PRECAUTIONS:

KEEP OUT OF REACH OF CHILDREN.

Avoid contact with skin and eyes. Wear protective gloves and coveralls when handling ROVRAL FLO.

Wash hands and exposed skin thoroughly after handling the concentrate and after application.

May be harmful if inhaled.

Avoid working in spray mist.

FIRST AID:

Remove patient from contaminated area, remove contaminated clothing. Keep patient warm, comfortable, at rest.

Eye contact: Irrigate with water for 15 minutes. Get medical attention.

Skin contact: Wash exposed skin with soap and water.

Ingestion: This product contains petroleum distillates. If ingested, DO NOT induce vomiting. If patient is conscious, wash out mouth. If breathing stops, start artificial respiration. Get medical attention.

Take container, label or product name and Pest Control Product Registration Number with you when seeking medical attention.

STORAGE CONDITIONS:

Protect from frost.

DISPOSAL:

1. Triple- or pressure-rinse the emptied container. Add the rinsings to the spray mixture in the tank.
2. Follow provincial instructions for any required additional cleaning of the container prior to its disposal.
3. Make the empty container unsuitable for further use.
4. Dispose of the container in accordance with provincial requirements.
5. For information on the disposal of unused, unwanted product, contact the manufacturer or the provincial regulatory agency. Contact the manufacturer and the provincial regulatory agency in case of a spill, and for clean-up of spills.

NOTICE TO USER: This control product is to be used only in accordance with the directions on the label. It is an offence under the Pest Control Products Act to use a control product under unsafe conditions.

NOTICE TO BUYER: Seller's guarantee shall be limited to the terms set out on the label and subject thereto, the buyer assumes the risk to persons or property arising from the use or handling of this product and accepts the product on that condition.

DIRECTIONS FOR USE:

For mixing, put one half the required water volume in the tank; stir in ROVRAL FLO and add the remainder of the water. Ensure mixing is complete. Agitate thoroughly before each application and continue to agitate during spraying. A prepared spray mixture should be used as soon as possible in the same day, especially if the water used for mixing is alkaline (greater than pH of 7). ROVRAL FLO should not be mixed with other pesticides, adjuvants or fertilizers except where stated.

CANOLA

CONTROL OF SCLEROTINIA STEM ROT

Rate: 2-3 L/ha ROVRAL FLO as a ground application in 100 litres of water per hectare, or as an aerial application in at least 45 litres of water per hectare. The use of the higher rate is recommended for fields with a history of heavy disease pressure or for dense crop stands. Good coverage of the plants is essential.

Timing: Apply ROVRAL FLO when the crop is in the 20-30% bloom stage. For Argentine varieties, the correct stage for application is when 15-20 flowers are open (including small pods) on the **main** stem. For Polish varieties, the correct stage for application is when 11-15 flowers (or small pods) are open on the **main** stem.

SUPPRESSION OF ALTERNARIA BLACK SPOT

Rate: 2 L/ha ROVRAL FLO as a ground application in 100 litres of water per hectare, or as an aerial application in at least 45 litres of water per hectare. Good coverage of the plants is essential. This disease mainly effects Polish (B. rapa) types of canola.

Aerial Applications:

Apply only by fixed-wing or rotary aircraft equipment which has been functionally and operationally calibrated for the atmospheric conditions of the area and the application rate and conditions of this label.

Label rates, conditions and precautions are product specific. Read and understand the entire label before opening this product. Apply only at the rate recommended for aerial

application on this label. Where no rate for aerial application appears for the specific use, this product cannot be applied by any type of aerial equipment.

Ensure uniform application. To avoid streaked, uneven or overlapped application, use appropriate marking devices.

Use Precautions:

Apply only when meteorological conditions at the treatment site allow for complete and even crop coverage. Apply only under conditions of good practice specific to aerial application as outlined in the *Basic Knowledge Requirements for Pesticide Education in Canada: Applicator Core* and *Aerial Module*, developed by CAPCO.

Do not apply to any body of water. Avoid drifting of spray onto any body of water or other non-target areas. Specified buffer zones should be observed.

Coarse sprays are less likely to drift, therefore, avoid combinations of pressure and nozzle type that will result in fine particles (mist). Do not apply during periods of dead calm or when wind velocity and direction pose a risk of spray drift. Do not spray when the wind is blowing towards a nearby sensitive crop, garden, terrestrial habitat (such as shelter-belt) or aquatic habitat.

Operator Precautions:

Do not allow the pilot to mix chemicals to be loaded onto the aircraft. Loading of premixed chemicals with a closed system is permitted.

It is desirable that the pilot have communication capabilities at each treatment site at the time of application.

The field crew and the mixer/loaders must wear chemical resistant gloves, coveralls and goggles or face shield during mixing/loading, cleanup and repair. Follow the more stringent label precautions in cases where the operator precautions exceed the generic label recommendations on the existing ground boom label.

All personnel on the job site must wash hands and face thoroughly before eating and drinking. Protective clothing, aircraft cockpit and vehicle cabs must be decontaminated regularly.

Product Specific Precautions:

Read and understand the entire label before opening this product. If you have questions, call the manufacturer at 1-888-AVENTIS (1-888-283-6847) or obtain technical advice from the distributor or your provincial agricultural representative. Application of this specific product must meet and/or conform to the following:

Volume: Apply the recommended rate in a minimum spray volume of 45 litres per hectare.

This label transcript service is offered by the Pest Management Regulatory Agency to provide efficient searching for label information. This service and this information do not replace the official hard-copy label. The PMRA does not provide any guarantee or assurance that the information obtained through this service is accurate, current or correct, and is therefore not liable for any loss resulting, directly or indirectly, from reliance upon this service.

+))