

1998/2000

DOMESTIC

500 g

C-I-L

EARWIG KILLER DUST

KEEP OUT OF REACH OF CHILDREN

READ THE LABEL BEFORE USING

GUARANTEE: chlorpyrifos 0.5%

REGISTRATION NO. 19759 P.C.P ACT

NU-GRO CORPORATION
Woodstock, Ontario N4S 8P6
Laval, Quebec H7S 1G3
Calgary, Alberta T2H 2S1

GENERAL INFORMATION

EARWIGS: Earwigs are brown, hard-shelled insects which have pincher-like appendages at the rear. These pinchers do not harm humans. Earwigs are nocturnal insects. They prefer dark damp areas. They feed on a variety of things such as flowers, leaves and vegetables. And they will also eat mites and the young stages of other insects. Food matter in homes is also attractive to them.

Earwigs overwinter as adults under rocks, logs, debris, in the soil, and around the foundation of the home. The female earwig lays her eggs in late winter to early spring brooding over them until they hatch. There are two generation of earwigs per year in most locations.

Control of earwigs is not simple. They are active insects and live in areas difficult to treat properly.

DIRECTION FOR USE

7.5 g = 1 level tablespoonful

OUTSIDE DWELLINGS: Soon after earwigs are first noticed, apply Earwig Killer Dust to the soil around the foundation of the home. Apply 500 g (full container) in 30cm band over a linear length of 40 m. Repeat the treatment when earwigs are again seen in the house or on the outside of the house. Make no more than 2 applications per season.

VEGETABLES: Apply dust to soil around plant at a rate of 70 g per 10 m of row. Use on broccoli, Brussels sprouts, cabbage, Chinese cabbage, cauliflower, carrots, celery, cucumbers, green peppers, onions, potatoes, sweet corn and tomatoes. Do not use on green bunching onions. Repeat as needed. Do not make more than nine applications during one season.

Days to wait between Crop
application and harvest

Crop

Broccoli	32
Brussels sprouts	32
Cabbage	32
Carrot	60
Chinese cabbage	32
Cauliflower	32
Celery	70
Cucumber	60
Green pepper	40
Onion	60
Potato	7
Sweet Corn	70
Tomato	70

ORNAMENTAL PLANTS & SHRUBS:

Soil Treatment: Dust turf or soil around plants to give even coverage. Apply 425 g per 10m². Make no more than 2 applications per season.

PRECAUTIONS: Avoid inhaling dust. Avoid contact with eyes, skin or clothing. May cause irritation to eyes or skin. Wash thoroughly after use. Avoid contamination of food, feed,

utensils, drinking water or any body of water. Wash contaminated food surfaces, utensils and clothing with soap and water. Keep children and pets away from treated areas.

DISPOSAL: Dispose of empty container in household garbage.

FIRST AID: If swallowed, induce vomiting by giving patient 1 to 2 glasses of water and inserting finger down throat. Obtain medical attention.

If on skin, wash with soap and water.

If in eyes, flush with water and obtain medical attention.

TOXICOLOGY: Chlorpyrifos is a cholinesterase inhibitor. Treat symptomatically. Atropine by injection only is an antidote.

NOTICE TO BUYER: Seller's guarantee shall be limited to the terms set out in the label and subject thereto, the buyer assumes the risk to persons or property arising from the use or handling of this product and accepts the product on that condition.

C-I-L is a trademark of ICI Canada Inc.
NU-GRO Corporation is a Registered User.

This label transcript service is offered by the Pest Management Regulatory Agency to provide efficient searching for label information. This service and this information do not replace the official hard-copy label. The PMRA does not provide any guarantee or assurance that the information obtained through this service is accurate, current or correct, and is therefore not liable for any loss resulting, directly or indirectly, from reliance upon this service.

+))